MARK SCHEME for the October/November 2013 series

9700 BIOLOGY

9700/41

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2013	9700	41

Mark scheme abbreviations

; R A AW <u>underline</u> max ora mp ecf I	separates marking points alternative answers for the same point reject accept (for answers correctly cued by the question, or by extra guidance) alternative wording (where responses vary more than usual) actual word given must be used by candidate (grammatical variants excepted) indicates the maximum number of marks that can be given or reverse argument marking point (with relevant number) error carried forward ignore
I	ignore
AVP	Alternative valid point (examples given as guidance)

Page 3		ge 3	Mark Scheme	Syllabus	Paper
			GCE AS/A LEVEL – October/November 2013	9700	41
1	(a)	allele – v	variation / different form, of a gene;		
		dominan	t – (allele) always expresses itself (in the phenotype wh	en present) ;	[2]
	(b)		ter the number of (CAG) repeats the earlier the sympton on a large time correlation ;	ns first appear /	inversely
		paired fig	gures;		[2]
	(c)	1. fear of	f needles ;		
		2. fear of	positive result ;		
		3. fear of	f effect of result on other members of family ;		
		4. no des	sire to have children ;		
		5. financi	ial / insurance, concerns / AW ;		
		6. possib	ility of false results ;		
		7. cost of	f test ;		
		8. not wo	orth having test because of no treatment ;		[max 3]
					[Total: 7]
2	(a)	in contex	t of woolly mammoth		
		1. individ	luals varied (in their phenotypes);		
		2. (pheno	otypic variation) caused by, genetic variation / mutation	;	
		3. chang	e in, selection pressure / environmental conditions;		
		4. <i>idea th</i>	nat variation increases the chance of some individuals s	urviving / AW;	
		5. nameo	d adaptation explained ; e.g. better insulation / smaller s	urface area to	volume
		6. survivo	ors breed ;		
		7. passe	d on <u>alleles</u> to offspring ;		

8. changed <u>allele</u> frequency (in population);

[max 5]

	Pa	ge 4	4		Mark Sc			Syllabus	Paper
				GCE AS/A L	EVEL – Oct	ober/Novembe	er 2013	9700	41
	(b)	1. c	difference	es in, primary	structure / se	equence of amir	no acids / p	olypeptide;	
		2. p	orovides	different, side	chains / R g	roups ;			
		3. c	change in	n, tertiary stru	cture / 3D sha	ape;			
		4. e	effect on	quaternary st	ructure ;				
		5. g	greater ef	ffect on β cha	in ;				
		6. c	change ir	n properties ;	A function				[max 3]
	(c)	(i)	1. still a	ble to offload	oxygen (in c	old temperature	es);		
			2. surfa	ice tissues co	lder than, cor	re / body, tempe	erature;		
			3. so ca	an maintain o	kygen supply	to surface tissu	ies;		[max 2]
		(ii)	1. no / t	iny, difference	e in effect of t	temperature on	haemoglob	oin alone ;	
			2. so no	o evidence (w	oolly mammo	oth haemoglobi	n) better ad	apted ;	
			-	ter reduction i ly mammoth		mperature on h	aemoglobir	n with red cell e	ffector in
			4. (so) v	woolly mamm	oth haemogle	obin (with red c	ell effector)	better adapted	to cold ;
			5. ref. c	hange to oxy	gen binding s	sites ;			
			6. so ca	an offload oxy	gen at low te	mperatures ;			[max 4]
									[Total: 14]
3	(a)	ade	enine / nit	trogen(ous) b	ase / purine ;	; R adenosine			
		ribo	ose / pen	tose ;					[2]
	(b)	1. (cell uses	s) ATP as sou	rce of energy	(;			
		2. /	ATP brok	en down ;					
		3. ((so) cell n	nust regenera	ate ATP;				
		4. f	rom ADF	P and Pi ;					
		5. r	ef. ADP	/ AMP, must I	be synthesise	ed in the cell;			[max 2]

Page 5				Syllabus	Paper	
				GCE AS/A LEVEL – October/November 2013	9700	41
	(c)	(i)	1. pa	almitic acid has more , hydrogens / C-H bonds ;		
			2. pe	er mole ;		
			3. hy	rdrogens needed for, ATP production / chemiosmosis / o	oxidative phosp	horylation; [max 2]
		(ii)	alani	ine – starvation / lack of fat or carbohydrate;		
			lacta	te – after anaerobic respiration ;		[2]
						[Total: 8]
4	(a)	(i)		ing ; e.g. 1st oestrogen peak at day 13, 2nd peak at day calculated number of days in between	y 41 / looked at	two peaks
			<u>28</u> ;			[2]
		(ii)	bega	an: day 13 or14 ;		
			ende	ed: day 29 or 30 ;		[2]
		(iii)	(ante	erior) pituitary (gland) ; R posterior pituitary		[1]
		(iv)	1. sti	mulates follicle;		
			2. to	secrete oestrogen ;		
			3. su	rge in LH secretion;		
			4. sti	imulates ovulation ;		
			5. re	f. development of corpus luteum / stimulates corpus lute	eum;	
			6. to	secrete progesterone;		[max 3]
	(b)	(i)	1. re	f. reliability ;		
			2. re	f. to irregularity of cycles ;		
			3. ide	<i>ea that</i> cannot be sure about menstrual phase on day 2	2;	
			4. ide	<i>ea that</i> using hormones alone might not identify day of o	cycle precisely e	enough ; [max 2]

	Page 6		6	Mark Scheme	Syllabus	Paper
				GCE AS/A LEVEL – October/November 2013	9700	41
		(ii)	1. (y	es because) oestrogen concentration high on day 22 an	d low on day 2	1
			2. (b	ut) shows correlation but not necessarily, linked / causa	l effect ;	
			3. co	oncentration of progesterone could be affecting performation	ance;	
			4. (p	rogesterone concentration) high at 22 days and low on o	day 2 ;	
			5. nc	ot LH as concentration low on both days ;		
			6. re	f. to small numbers in investigation / more evidence nee	eded ;	
			7. re	f. to use of statistics to determine if difference in results	is significant ;	[max 4]
						[Total: 14]
5	(a)	1. r	no cha	ange between 1860 and 1930 ;		
		2. r	ef. to	increases from 1930 to 2010;		
		3. ι	use of	figures including <u>units</u> ;		[3]
	(b)	1. s	single-	-cross hybrids have homozygous parents;		
		2. e	each h	nas inherited the same alleles ;		
		3. (so) th	ey are uniformly heterozygous;		
		4. c	double	e-cross hybrids have heterozygous parents ;		
				nas inherited different combinations of alleles		
		-	or mixtui	re of) homozygous dominant, homozygous recessive a	nd heterozygou	s hybrids ; [max 3]
	(c)	(i)	1. th	e greater the inbreeding coefficient, the lower the yield ;		
			2. in	each site in each year ;		
			3. us	se of figures ;		[max 2]
		(ii)	1. th	e yield differs, at different sites / in different years ;		
			2. fo	r the same inbreeding coefficient;		
			3. us	se of figures ;		
			4. na	amed environmental factor; e.g. rainfall / temperature /	mineral content	of soil [max 2]
						[Total: 10]
						-

	Pa	ge 7	,	Mark Scheme	Syllabus	Paper
	(-)	(1)		GCE AS/A LEVEL – October/November 2013	9700	41
6	(a)	(i)	-	t <u>er</u> speed (if myelinated) ;		[0]
		/::)		parative figures with units ;		[2]
		(ii)	-	er diameter greater speed / ora ;		[0]
			com	parative figures with units ;		[2]
	(b)	1. r	nyelin	insulates <u>axon</u> ;		
		2. r	no my	elin at nodes ;		
		3. a	action	potentials / depolarisation, only at nodes (of Ranvier);		
		4. le	ocal c	ircuits set up between nodes ;		
		5. a	action	potentials 'jump' from node to node / saltatory conduction	on;	
		6. r	nyelin	ation prevents leakage of ions ; ora		[max 3]
	(c)	(i)	1. (s	heath) treated as, 'foreign' / non-self ;		
			2. re	f. role of, antibodies / phagocytes / lymphocytes ;		[2]
		(ii)	1. le:	ss insulation of <u>axon</u> ;		
			2. ac	ction potentials, slow down / stop ;		[2]
						[Total: 11]
7	(a)	(i)	1. (b	lue) light is absorbed and used for photosynthesis;		
			2. C	O_2 , used / concentration decreased ;		
			3. lea	ads to, rise in pH / decrease in acidity ;		[max 2]
		(ii)	1. re	spiration but no photosynthesis ;		
			2. C	O ₂ , produced / released ;		
			3. lea	ads to, decrease in pH / increase in acidity ;		[max 2]
	(b)	(i)	absc	orb light (energy) ;		
			pass	; (light) <u>energy</u> onto, primary pigment / chlorophyll a / rea	action centre;	[2]
		(ii)	H₂O	$\rightarrow 2H^+ + 2e^- + \frac{1}{2}O_2;$		
			A 2F	$H_2O \longrightarrow 4H^+ + 4e^- + O_2$		[1]
	((iii)		a / thylakoid, <u>membrane</u> ;		[1]
	·	-				[Total: 8]

	Page 8		Mark Scheme	Syllabus	Paper
			GCE AS/A LEVEL – October/November 2013	9700	41
8	(a)	any num	ber between 873 – 882 inclusive ;;		
		allow on	e mark for correct working or for number not rounded up		[max 2]
	(b)	named s	pecies (no mark)		
		e.g. anin direct hu habitat d climate d increase spread / lack of fc increase e.g. plan direct hu habitat d climate c increase spread / loss of p	<pre>vant reasons for a named species ;;;; nal species man effect e.g. hunting / fishing / collection / skins estruction change qualified in pollution increase, in disease or new disease ood d predation t species man effect e.g. specimen collection / logging estruction change qualified in pollution increase, in disease or new disease ollinators d competition from introduced plants</pre>		[4]
					[Total: 6]
9	dor	mancy;			
	eml	oryo ;			
	aleı	urone;			
	end	losperm ;			
	mal	tose ;			
	ATF	⊃ / energy	′;		
	trar	scription	/ expression ;		[7]

 10 (a) 1. chance / random / spontaneous ; 2. change in, base / nucleotide, sequence (in DNA) ; 3. during DNA replication ; 4. base substitution ; 5. often no effect / silent mutation / may code for same amino acid ; 6. base addition / base deletion ; 7. have great effect on phenotype ; 8. frame shifts ; 9. alters whole sequence of bases after mutation ; 10. may lead to stop codon ; 11. different / new, <u>allele</u> ; 12. protein, different shape / different function / not made ; (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; 		Pa	ge 9	Mark Scheme	Syllabus	Paper
 2. change in, base / nucleotide, sequence (in DNA); 3. during DNA replication; 4. base substitution; 5. often no effect / silent mutation / may code for same amino acid; 6. base addition / base deletion; 7. have great effect on phenotype; 8. frame shifts; 9. alters whole sequence of bases after mutation; 10. may lead to stop codon; 11. different / new, <u>allele;</u> 12. protein, different shape / different function / not made; (b) 1. no / no functional, channels for Cl⁻ ions; 2. Cl⁻ ions do not move out; 3. less water leaves cell; 4. mucus (on cell surface membrane) stays, thick / sticky; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 				GCE AS/A LEVEL – October/November 2013	9700	41
 3. during DNA replication ; 4. base substitution ; 5. often no effect / silent mutation / may code for same amino acid ; 6. base addition / base deletion ; 7. have great effect on phenotype ; 8. frame shifts ; 9. alters whole sequence of bases after mutation ; 10. may lead to stop codon ; 11. different / new, <u>allele</u> ; 12. protein, different shape / different function / not made ; (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; 	10	(a)	1. chanc	e / random / spontaneous ;		
 4. base substitution ; 5. often no effect / silent mutation / may code for same amino acid ; 6. base addition / base deletion ; 7. have great effect on phenotype ; 8. frame shifts ; 9. alters whole sequence of bases after mutation ; 10. may lead to stop codon ; 11. different / new, <u>allele</u> ; 12. protein, different shape / different function / not made ; (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 			2. chang	e in, base / nucleotide, sequence (in DNA) ;		
 5. often no effect / silent mutation / may code for same amino acid; 6. base addition / base deletion; 7. have great effect on phenotype; 8. frame shifts; 9. alters whole sequence of bases after mutation; 10. may lead to stop codon; 11. different / new, <u>allele;</u> 12. protein, different shape / different function / not made; (b) 1. no / no functional, channels for Cl⁻ ions; 2. Cl⁻ ions do not move out; 3. less water leaves cell; 4. mucus (on cell surface membrane) stays, thick / sticky; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 11. blocked pancreatic duct; 			3. during	DNA replication ;		
 6. base addition / base deletion; 7. have great effect on phenotype; 8. frame shifts; 9. alters whole sequence of bases after mutation; 10. may lead to stop codon; 11. different / new, <u>allele</u>; 12. protein, different shape / different function / not made; (b) 1. no / no functional, channels for Cl⁻ ions; 2. Cl⁻ ions do not move out; 3. less water leaves cell; 4. mucus (on cell surface membrane) stays, thick / sticky; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 11. blocked pancreatic duct; 			4. base s	substitution;		
 7. have great effect on phenotype ; 8. frame shifts ; 9. alters whole sequence of bases after mutation ; 10. may lead to stop codon ; 11. different / new, <u>allele</u> ; 12. protein, different shape / different function / not made ; [m (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; [m 			5. often r	no effect / silent mutation / may code for same amino a	cid;	
 8. frame shifts; 9. alters whole sequence of bases after mutation; 10. may lead to stop codon; 11. different / new, <u>altele</u>; 12. protein, different shape / different function / not made; (b) 1. no / no functional, channels for Cl⁻ ions; 2. Cl⁻ ions do not move out; 3. less water leaves cell; 4. mucus (on cell surface membrane) stays, thick / sticky; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 11. blocked pancreatic duct; 			6. base a	addition / base deletion;		
 9. alters whole sequence of bases after mutation ; 10. may lead to stop codon ; 11. different / new, <u>allele</u>; 12. protein, different shape / different function / not made ; [m (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 			7. have g	great effect on phenotype ;		
 10. may lead to stop codon ; 11. different / new, <u>allele</u>; 12. protein, different shape / different function / not made ; [m (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; [m 			8. frame	shifts ;		
 11. different / new, <u>allele</u>; 12. protein, different shape / different function / not made; [m (b) 1. no / no functional, channels for Cl⁻ ions; 2. Cl⁻ ions do not move out; 3. less water leaves cell; 4. mucus (on cell surface membrane) stays, thick / sticky; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 11. blocked pancreatic duct; [m 			9. alters	whole sequence of bases after mutation;		
 12. protein, different shape / different function / not made ; [m (b) 1. no / no functional, channels for Cl⁻ ions ; 2. Cl⁻ ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; [m 			10. may	lead to stop codon ;		
 (b) 1. no / no functional, channels for Cl' ions; 2. Cl' ions do not move out; 3. less water leaves cell; 4. mucus (on cell surface membrane) stays, thick / sticky; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 11. blocked pancreatic duct; 			11. differ	rent / new, <u>allele</u> ;		
 2. Cl' ions do not move out ; 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 			12. prote	ein, different shape / different function / not made ;		[max 9]
 3. less water leaves cell ; 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 		(b)	1. no / no	o functional, channels for Cl ⁻ ions ;		
 4. mucus (on cell surface membrane) stays, thick / sticky ; 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 			2. Cl⁻ ion	is do not move out ;		
 5. symptoms – any 4 from: mucus not moved effectively by cilia / mucus accumulates; 6. reduced gaseous exchange / longer diffusion pathway; 7. difficulty in breathing; 8. more infections / (mucus) traps bacteria; 9. lungs are scarred; 10. blocked sperm ducts; 11. blocked pancreatic duct; [m 			3. less w	vater leaves cell;		
 mucus not moved effectively by cilia / mucus accumulates ; 6. reduced gaseous exchange / longer diffusion pathway ; 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 			4. mucus	s (on cell surface membrane) stays, thick / sticky ;		
 7. difficulty in breathing ; 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 						
 8. more infections / (mucus) traps bacteria ; 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 			6. reduce	ed gaseous exchange / longer diffusion pathway;		
 9. lungs are scarred ; 10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m 			7. difficul	Ity in breathing;		
10. blocked sperm ducts ; 11. blocked pancreatic duct ; [m			8. more i	infections / (mucus) traps bacteria ;		
11. blocked pancreatic duct ; [m			9. lungs	are scarred;		
			10. block	ked sperm ducts ;		
[Total			11. block	ked pancreatic duct ;		[max.6]
						[Total: 15]

Paç	ge 10	Mark Scheme	Syllabus	Paper
1 (a)	1 poultic	GCE AS/A LEVEL – October/November 2013	9700	41
1 (a)	1. multic			
		are) differentiated into tissues ;		
		rophic / photosynthetic ;		
		yotic (cells);		
		n is storage compound ;		
		e have) chloroplasts / chlorophyll ;		
	7. cell w	all;		
	8. made	of cellulose;		
	9. plasm	nodesmata ;		
	10. large	e (central) vacuole ;		[max 7
(b)	1. 0.5–1	.0 μ m, diameter / width ;		
	2. doubl	e membrane ;		
	3. inner	membrane folded / cristae ;		
	4. hold,	stalked particles / ATP synthase / ATP synthetase;		
	5. site of	f ETC ;		
	6. ref. H	$^{\scriptscriptstyle +}$ and intermembrane space ;		
	7. ATP p	production;		
	8. oxida	tive phosphorylation / chemiosmosis;		
	9. matrix	is site of, link reaction / Krebs cycle ;		
	10. enzy	vmes in matrix ;		
	11. 70S	ribosomes ;		
	12. (mito	ochondrial) DNA ;		[max 8

[Total: 15]