
CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

May/June 2016

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.


Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
1 (a) (i)	dative (covalent) <i>or</i> coordinate Hydrogen/H (bonding)	2
(ii)	octahedral	1
(iii)	$\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O} \rightarrow \text{Mg}(\text{NO}_3)_2 + 6\text{H}_2\text{O}$ $\text{Mg}(\text{NO}_3)_2 \rightarrow \text{MgO} + 2\text{NO}_2 + \frac{1}{2}\text{O}_2$ <p><i>any three of</i> (solid) dissolves/turns to liquid condensation on tube <u>white</u> solid (forms/remains) brown fumes (evolved) gas formed that relights a glowing splint</p>	4
(iv)	M_r values: $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O} = 256.3$ $\text{MgO} = 40.3$ or (loss in molar mass = $256.3 - 40.3 = 216$) percentage loss = $100 \times 216 / 256.3 = \mathbf{84.3 / 84.4\%}$	2
(b)	(cat)-ionic radius / ion size increases (down the group) less polarisation / distortion of nitrate ion / NO_3^-	2
(c)	$2\text{AgNO}_3 \rightarrow 2\text{Ag} + 2\text{NO}_2 + \text{O}_2$	1
		[Total: 12]
2 (a) (i)	(an acid that is) partially / incompletely ionised / dissociated	1
(b) (i)	$\text{p}K_a = -\log K_a$ or $K_a = 10^{-\text{p}K_a}$	1



Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
(ii)	<p>ethanoic acid (1) is more acidic than propanoic acid (2) due to smaller electron-donating (R/ alkyl) group/ less electron-donating (R/ alkyl) group(s)</p> <p>2-chloropropanoic acid (3) is more acidic than propanoic acid (2) due to electron-withdrawing/ electronegative (Cl/ chlorine) atom</p> <p>2-chloropropanoic acid (3) is more acidic than 3-chloropropanoic acid (4) since the Cl/ chlorine/ electronegative atom is closer to the CO_2^-/ acid</p>	3
(c) (i)	 <p>M1: voltmeter / V and salt bridge labelled</p> <p>M2: Cu and Cu^{2+} / CuSO_4 (any soluble Cu(II) salt)</p> <p>M3: H_2 (arrow in) and H^+ / HCl / H_2SO_4 / any mineral acid</p> <p>M4 Pt and one solution at 1 M / 1 mol dm^{-3} OR H_2 at 1 atm</p>	4
(ii)	<p>$E^\ominus_{\text{cell}} = 0.34 \text{ (V)}$ and $(\text{Cu}^{2+})/\text{Cu}$ is the positive electrode</p>	1
d (i)	<p>$K_a = 1.23 \times 10^{-5}$</p> <p>$[\text{H}^+] = \sqrt{K_a \cdot c} = \sqrt{(1.23 \times 10^{-5} \times 0.1)} = 1.11 \times 10^{-3} \text{ mol dm}^{-3}$</p> <p>pH = 3.0 (2.96) ecf from $[\text{H}^+]$</p>	2


Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
(ii)	$E = 0.0 + 0.059\log(1.11 \times 10^{-3})$ OR $= -0.17(4)V$ so new $E_{\text{cell}} = 0.34 + 0.17 = \mathbf{0.51V}$ ecf from (d)(i)	2
		[Total: 14]
3 (a) (i)	$(\text{CH}_3)_2\text{CHCN}$	1
(ii)	reaction 1: NH_3 (in ethanol) under pressure (+ heat) or heat NH_3 in a sealed tube reaction 2: KCN/NaCN and heat/reflux (in ethanol) reaction 3: $\text{H}_2 + \text{Ni}$ or LiAlH_4	3
(b) (i)	$\text{CH}_3\text{CH}_2\text{NH}_2 + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{CH}_2\text{NH}_3^+ (+) \text{OH}^-$	1
(ii)	ethylamine is more basic than ammonia... because of electron-donating (alkyl/ethyl/R) group (in ethylamine) which makes the <u>lone pair</u> (on N) more available for donation or the <u>lone pair</u> (on N) more available for a proton/ H^+	2
(c) (i)	A solution which resists/minimises/roughly maintains changes in <u>pH</u> when (small amounts of) H^+ or OH^- are added	1
(ii)	$\text{CH}_3\text{NH}_2 + \text{H}^+ \rightarrow \text{CH}_3\text{NH}_3^+$ $\text{CH}_3\text{NH}_3\text{Cl} + \text{OH}^- \rightarrow \text{CH}_3\text{NH}_2 + \text{H}_2\text{O} + \text{Cl}^-$	2
		[Total: 10]


Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
4 (a) (i)	 <p>(cis) (trans)</p>	2
(ii)	<p>cis is (more) polar due to both Cl^(δ-) on same side or cis is (more) polar as dipoles do not cancel / unsymmetrical or trans is non-polar as its bond dipoles cancel</p>	1
(iii)	<p>(This can only be <i>cis</i>) its mirror image is the same / superimposable</p> <p>or the distance between two coordinating nitrogens / oxygens is too small to bond <i>trans</i> or difficult for the NH₂ and O to change places (since 5-membered rings can only bridge adjacent positions)</p>	1
(b) (i)	It's not square planar or it's tetrahedral	1
(ii)	<p>must be 3D structure (i.e. tetrahedral-like)</p>  <p><i>or</i> <i>etc</i></p>	1
		[Total: 6]


Page 7	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
6 (a)	<p><i>essential mark</i></p> <p>M1 the reactants/substrate has a shape complementary/specific to active site – can be awarded from a labelled diagram as below or diagrams showing this specificity clearly</p> <p><i>any two of</i></p> <p>M2: reactants/substrate binds to/fits into the active site of the enzyme</p> <p>M3: (Interaction with site) causes a specific bond to be weakened, (which breaks) or lowers activation energy</p> <p>M4: forms an E-S complex</p> <p>M5: products released from enzyme/active site</p> <p>labelled diagrams</p> 	3
(b) (i)	<p>δ 26 is CH₃-CO δ 52 is CH₃-O</p> <p>δ 169 is CH₃CO δ 167 is phenyl-CO</p> <p><u>Phenyl ethanoate</u> is B <u>methyl benzoate</u> is A</p> <p>M1 = any two correct δ linked to phenylethanoate/methyl benzoate</p> <p>M2 = the rest correct</p>	2
(ii)	<p>heat with H₃O⁺ (to hydrolyse the ester)</p> <p>then add Br₂(aq)/bromine water</p> <p>decolourises/gives white ppt. (with phenol from B)</p>	3
		[Total: 8]



Page 8	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
7 (a) (i)	<p>labelled with</p> <p>M1: <u>DC</u> power supply + and – / battery / cell / + and – sign (on cell / electrodes) with a complete circuit</p> <p>M2: buffer solution / electrolyte labelled</p> <p>M3: (amino acid) mixture / x on (filter) paper / gel / agarose</p> 	3
(ii)	<p>direction of movement related to charge (of amino acids)</p> <p>distance travelled depends on charge / M_r (of amino acids)</p>	2
(b) (i)	<p>Asp + Val:</p> <p>pH 12 because Asp will be $-\text{CH}_2\text{COO}^-$ (R-group) moves further (to positive electrode than Val)</p> <p>or pH 12 Asp more negative so moves further (to positive electrode)</p> <p>or pH 12 because Asp has a charge of 2– but Val has a charge of 1–</p> <p>or best at pH 7 because Asp will be negatively charged (anionic) but Val neutral</p>	1
(ii)	<p>Lys + Ser:</p> <p>pH 2 because Lys will be $(\text{CH}_2)_4\text{NH}_3^+$ (R-group) moves further (to negative electrode than Ser)</p> <p>or pH 2 Lys more positive so moves further (to negative electrode)</p> <p>or pH 2 because Lys has a charge of 2+ and Ser has a charge of 1+</p> <p>or pH 7 because Lys is positively charged (cationic) but Ser neutral / zwitterionic</p>	1




Page 9	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
(iii)	Tyr + Phe: pH 12 because Tyr will be $\text{C}_6\text{H}_5\text{CH}_2\text{O}^-$ (R-group) moves further / more / faster (to positive electrode than Phe) or pH12 because Tyr has a charge of 2- but Phe has a charge of 1-	1
(c) (i)	 <p>M1: for $-\text{CONH}-$ as shown above</p> <p>M2: for rest of molecule and correct connectivity of the bonds</p>	2
(ii)	<i>from the IR spectrum</i> <ul style="list-style-type: none"> • E is O-H or N-H (allow NH_2) • F is C=O • G is C-O 	2
		[Total: 12]
8 (a)	<p>M1: solubility increases (down the group)</p> <p>M2: because lattice energy decreases faster than does ΔH_{hyd}</p> <p>M3: ΔH_{sol} / enthalpy of solution becomes more exothermic / less endothermic</p>	3
(b) (i)	Should be the same / similar (enthalpy change), as (both acids) are fully ionised / strong acids	1

Page 10	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
(ii)	$\text{Ca(s)} + 2\text{H}^+(\text{aq}) \longrightarrow \text{Ca}^{2+}(\text{aq}) + \text{H}_2(\text{g})$  <p>gas phase ions: $\text{Ca}^{2+}(\text{g}) + 2\text{H}^+(\text{g})$</p> $x = \Delta H_{\text{at}}(\text{Ca}) + \text{IE}(1) + \text{IE}(2) - 2\Delta H_{\text{hyd}}(\text{H}^+) + \Delta H_{\text{hyd}}(\text{Ca}^{2+}) - 2\text{IE}(\text{H}) - E(\text{H}-\text{H})$ $x = 178 + 590 + 1150 + 2(1090) - 1576 - 2(1310) - 436$ $x = -534 \text{ kJ mol}^{-1}$	4
(c)	<p>$\text{CH}_3\text{CO}_2\text{H}$ is incompletely ionised / weak acid / weaker acid</p> <p>enthalpy change of ionisation (of CH_3COOH) is $+2 \text{ kJ mol}^{-1}$</p> <p>or energy needed to ionise / dissociate (CH_3COOH)</p>	2
		[Total: 10]
9 (a)		1

Page 11	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9701	42

Question	Answer	Marks
(b)	<p>H is </p> <p>J is </p> <p>or </p> <p style="text-align: center;">J1 J2</p>	2
(c)	<p>step 1: $(\text{CH}_3)_2\text{CHCH}_2\text{Cl} + \text{AlCl}_3$ (+ heat)</p> <p>step 2: $\text{CH}_3\text{COCl} + \text{AlCl}_3$ (+ heat)</p> <p>step 3: $\text{HCN} + \text{NaCN}$ <i>or</i> $\text{HCN} + \text{base}$ <i>or</i> $\text{HCN} + \text{CN}^-$</p> <p><i>(steps 4 and 5 could be reversed on J)</i> If J1 step 4 then step 5 J2 step 5 then step 4</p> <p>step 4: H_3O^+ + heat/aqueous HCl + heat</p> <p>step 5: conc H_2SO_4 + heat/ conc H_3PO_4 + heat <i>or</i> Al_2O_3 + heat</p> <p>step 6: $\text{H}_2 + \text{Ni}$ (+ heat)</p>	6
(d)	<p>step 1: electrophilic substitution <i>or</i> alkylation</p> <p>step 6: reduction / hydrogenation / addition</p>	2
		[Total: 11]

