Cambridge International AS & A Level

89702

N

Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME		
CENTRE NUMBER		CANDIDATE NUMBER
CHEMISTRY 9701/22		
Paper 2 AS Level Structured Questions		October/November 2016
		1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units. A Data Booklet is provided.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **10** printed pages and **2** blank pages.



Answer **all** the questions in the spaces provided.

1 A 0.50g sample of a Group 2 metal, **M**, was added to 40.0 cm³ of 1.00 mol dm⁻³ hydrochloric acid (an excess).

equation 1 $M(s) + 2HCl(aq) \rightarrow MCl_2(aq) + H_2(g)$

(a) Calculate the amount, in moles, of hydrochloric acid present in 40.0 cm³ of 1.00 mol dm⁻³ HC*l*.

amount = mol [1]

(b) When the reaction had finished, the resulting solution was made up to 100 cm³ in a volumetric flask.

A 10.0 cm³ sample of the solution from the volumetric flask required 15.0 cm^3 of $0.050 \text{ mol dm}^{-3}$ sodium carbonate solution, Na_2CO_3 , for complete neutralisation of the remaining hydrochloric acid.

(i) Write the equation for the complete reaction of sodium carbonate with hydrochloric acid.

......[1]

(ii) Calculate the amount, in moles, of sodium carbonate needed to react with the hydrochloric acid in the 10.0 cm³ sample from the volumetric flask.

amount = mol [1]

(iii) Calculate the amount, in moles, of hydrochloric acid in the 10.0 cm³ sample.

amount = mol [1]

(iv) Calculate the total amount, in moles, of hydrochloric acid remaining after the reaction shown in equation 1.

amount = mol [1]

(v) Use your answers to (a) and (b)(iv) to calculate the amount, in moles, of hydrochloric acid that reacted with the 0.50 g sample of M.

amount = mol [1]

(vi) Use your answer to (v) and equation 1 to calculate the amount, in moles, of **M** in the 0.50g sample.

amount = mol [1]

(vii) Calculate the relative atomic mass, A_r , of **M** and identify **M**.

 $A_{\rm r}$ of **M** =

identity of **M** =

[2]

[Total: 9]

2 Dinitrogen tetraoxide, N₂O₄, and nitrogen dioxide, NO₂, exist in dynamic equilibrium with each other.

 $N_2O_4(g) \rightleftharpoons 2NO_2(g) \qquad \Delta H = +54 \text{ kJ mol}^{-1}$

The energy profile for this reaction is shown.



- (a) Add labelled arrows to the energy profile to indicate
 - the enthalpy change of the reaction, ΔH ,
 - the activation energy of the forward reaction, E_a .

[2]

- (b) $0.0500 \text{ mol of } N_2O_4$ was placed in a sealed vessel of volume 1.00 dm^3 , at a temperature of $50 \degree \text{C}$ and a pressure of $1.68 \times 10^5 \ \text{Pa}$. The mass of the resulting equilibrium mixture was $4.606 \ \text{g}$.
 - (i) Calculate the average molecular mass, M_r , of the resulting equilibrium mixture. Give your answer to **three** significant figures.

*M*_r = [2]

(ii) The number of moles of N_2O_4 that dissociated can be represented by *n*.

State, in terms of n, the amount, in moles, of NO₂ in the equilibrium mixture.

moles of $NO_2 = \dots$ [1]

The number of moles of N_2O_4 remaining at equilibrium is (0.05 - n).

(iii) State, in terms of *n*, the total amount, in moles, of gas in the equilibrium mixture.

[1]

(iv) State, in terms of n, the mole fraction of NO₂ in the equilibrium mixture.

[1]

In this equilibrium mixture, the mole fraction of NO_2 is 0.400.

(v) Use your answers to (ii) and (iv) to calculate the amount in moles of each gas in the equilibrium mixture. Give your answers to **three** significant figures.

amount of $N_2O_4 = \dots$	mol
amount of NO ₂ =	mol [2]

(vi) Write the expression for the equilibrium constant, K_{p} , for this equilibrium.

 $K_{p} =$

[1]

(vii) Use the total pressure of the mixture, 1.68×10^5 Pa, to calculate the value of the equilibrium constant, K_p , and give its units.

*K*_p =

units =[3]

[Total: 13]

[Turn over

- 3 The Periodic Table is arranged such that the properties of the elements show a number of trends.
 - (a) A plot of the first ionisation energies for the first 18 elements is shown.



(b) A plot of the melting points of the elements across the third period is shown.



7

(i) Explain the increase in melting point from atomic number 11 to 12.

(ii)	Suggest a reason why the increase from atomic number 12 to 13 is much smaller than the increase from atomic number 11 to 12.
(iii)	State and explain the pattern of the melting points from atomic number 15 to 18.
	[3]
(iv)	Explain why the element with atomic number 14 has a melting point so much higher than the rest of the elements in the third period.
	[1]
	[Total: 15]

[Turn over

- 4 In each section of this question the structural formula of an organic compound is shown. For each compound answer the questions about it.
 - (a) CH₃CH₂CHBrCH₃
 - (i) Name this compound.
 -[1]
 - (ii) This compound shows stereoisomerism.

Draw the two stereoisomers in the conventional way.

(iii) Give the structures of **three** other structural isomers of C_4H_0Br .



[2]

(b) $(C_2H_5)_3CBr$

(i) Name this compound.

(ii) $(C_2H_5)_3CBr$ reacts with aqueous OH⁻.

Complete the mechanism for this reaction including all necessary curly arrows, charges, partial charges and lone pairs.

 (c) $CH_3CH_2CH_2CHBrCH_3$

(i) Give the reagents and conditions necessary for the conversion of this compound into a mixture of alkenes.



(iii) Draw the skeletal formulae of the three alkenes produced by the conversion in (i).







[3]

[Total: 17]

(a) Six particles are listed. H• Cl• Cl-H⁺ •CH₃ ⁺CH₃ (i) Identify two particles produced during the reaction of methane and chlorine in the presence of UV light.[1] (ii) Identify the **two** particles produced by the heterolytic fission of a bond in chloromethane. (b) Seven reaction types are listed. addition substitution oxidation elimination hydrolysis condensation reduction (i) Name the type of reaction involved when Tollens' reagent is used to identify an aldehyde. (ii) Name the type of reaction involved in the test for a carbonyl group using 2,4-DNPH. (iii) Name the type of reaction involved in the reaction of a ketone with NaBH₄.[1] (iv) Name the type of reaction involved in the reaction of an aldehyde with HCN.

......[1]

[Total: 6]

In each section of this question choose the answer or answers from the options listed.

5

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.