

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/31

Paper 3 Advanced Practical Skills 1

May/June 2017

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Give details of the practical session and laboratory where appropriate, in the boxes provided.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.
Electronic calculators may be used.
You may lose marks if you do not show your working or if you do not use appropriate units.
Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.
A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [] at the end of each question or part question.

Session	
Laboratory	

For Examiner's Use	
1	
2	
3	
Total	

This document consists of **12** printed pages.



- 1 In this experiment you will determine the relative formula mass of a copper salt by titration.

A solution of the copper salt reacts with excess acidified potassium iodide, producing iodine. This iodine is then titrated with aqueous sodium thiosulfate, using starch indicator.

FA 1 is an aqueous solution of the copper salt prepared by dissolving 26.0g of the salt to make 1.00 dm³ of solution.

FA 2 is dilute sulfuric acid, H₂SO₄.

FA 3 is aqueous potassium iodide, KI.

FA 4 is 0.110 mol dm⁻³ sodium thiosulfate, Na₂S₂O₃.
starch indicator

(a) Method

- Fill the burette with **FA 4**.
- Pipette 25.0 cm³ of **FA 1** into a conical flask.
- Use the measuring cylinder to add approximately 10 cm³ of **FA 2** to the same conical flask.
- Use the measuring cylinder to add approximately 20 cm³ of **FA 3** to the mixture in the conical flask. The mixture will now be a brown colour, due to iodine produced in the reaction.
- Begin your rough titration by adding **FA 4** from the burette until the mixture becomes light brown.
- Add 10 drops of starch indicator. The mixture will become darker.
- Continue titrating until the mixture becomes an off-white colour. This is the end-point.
- Add **one** drop of starch indicator to check that no traces of dark colour are produced. If the mixture stays off-white, the titration is finished. If some dark colour is produced, because iodine is still present, continue the titration.
- Record your burette readings and the rough titre in the space below.

The rough titre is cm³.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of **FA 4** added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	

Keep FA 3 and starch indicator for use in Question 3.

[7]

- (b)** From your accurate titration results, obtain a suitable value for the volume of **FA 4** to be used in your calculations.
Show clearly how you obtained this value.

The iodine produced required cm³ of **FA 4**. [1]


(c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- (i)** Calculate the number of moles of sodium thiosulfate, $\text{Na}_2\text{S}_2\text{O}_3$, in the volume of **FA 4** calculated in **(b)**.

moles of $\text{Na}_2\text{S}_2\text{O}_3$ = mol


- (ii)** Balance the equation for the reaction of iodine with sodium thiosulfate. State symbols are not required.



- (iii)** Using your answer to **(ii)**, calculate the number of moles of iodine that reacted with the number of moles of $\text{Na}_2\text{S}_2\text{O}_3$ calculated in **(i)**.

moles of I_2 = mol

- (iv)** Iodine, I_2 , is produced in the reaction between **FA 1** and **FA 3**. **FA 3** is in excess.



Using your answer to **(iii)**, calculate the number of moles of copper(II) ions in 25.0 cm^3 of **FA 1**.

moles of Cu^{2+} ions = mol

- (v)** Using your answer to **(iv)** and the information on page 2, calculate the relative formula mass of the copper compound in **FA 1**.

M_r of copper compound =
[4]

[Total: 12]

- 2 Malachite is a basic form of copper carbonate in which copper hydroxide is also present. The accepted chemical formula of malachite is $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$.

When malachite is heated, it decomposes as shown.



In this experiment, you will heat malachite to decompose it and use your results to obtain evidence about the accepted formula of malachite.

FA 5 is malachite, $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$.

(a) Method

Read through the method before starting any practical work.

In the space below prepare a **single** table for your results of **Experiments 1** and **2**.

Experiment 1

- Weigh a crucible with its lid and record the mass.
- Add between 2.5g and 3.0g of **FA 5** to the crucible. Weigh the crucible with **FA 5** and lid and record the mass.
- Place the crucible on the pipe-clay triangle.
- Heat the crucible and contents gently for about two minutes, with the lid on.
- Remove the lid and continue heating gently for about three minutes.
- Replace the lid and leave the crucible and residue to cool for at least five minutes. Then reweigh the crucible and contents with the lid on. Record the mass.
- **While the crucible is cooling, you may wish to begin work on Question 3.**
- Calculate and record the mass of **FA 5** used and the mass of residue obtained.
- State the observation(s) you made while the reaction was taking place.

.....

Experiment 2

Repeat the method used in **Experiment 1**, using between 1.5g and 2.0g of **FA 5** in the second crucible.

Results

I	
II	
III	
IV	
V	
VI	

[6]

(b) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- (i)** Use your results from **Experiment 1** to calculate the number of moles of copper oxide, CuO, obtained as residue.
Use the Periodic Table on page 12 for any data you may require.

moles of CuO obtained in **Experiment 1** = mol

- (ii)** Use your answer to **(i)**, the equation on page 4 and the mass of **FA 5** you used in **Experiment 1**, to calculate the relative formula mass, M_r , of malachite.

M_r of malachite (from **Experiment 1**) =

- (iii)** Use your results from **Experiment 2** to calculate another value for the relative formula mass, M_r , of malachite.

M_r of malachite (from **Experiment 2**) =

- (iv)** Use data from the Periodic Table to calculate the relative formula mass, M_r , of malachite from its accepted formula, $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$.

M_r of malachite (from formula) =

- (v)** If the relative formula mass of malachite obtained from **either** of your experiments is within 2.5% of the answer in **(iv)**, this is good evidence that the accepted formula, $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$, is correct.

Show by calculation whether either of your experiments supports the accepted formula.

[5]

(c) (i) State **one** way of improving the accuracy of the experimental method, using the same masses of **FA 5**.

Explain the benefit of your improvement.

.....

.....

.....

.....

(ii) Explain why you would expect **Experiment 1** to be more accurate than **Experiment 2**.

.....

.....

.....

[3]

[Total: 14]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where reagents are selected for use in a test, the **name** or **correct formula** of the element or compound must be given.

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

(a) **FA 6** is another salt of copper. The anion present is one of those listed in the Qualitative Analysis Notes.

- (i) Transfer a **small** spatula measure of **FA 6** into a hard-glass test-tube. Heat gently at first, then heat strongly, until no further change occurs.

Record **all** your observations below.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Suggest the chemical formula of **FA 6**.

.....

[3]

- (b) (i)** Dissolve the remainder of **FA 6** in an approximately 10 cm depth of distilled water in a boiling tube.

FA 7 is a solution of a salt containing one anion from those listed in the Qualitative Analysis Notes.

Two cations are also present.

Carry out the tests described below using separate portions of solutions **FA 6** and **FA 7**. Record your observations in the table.

<i>test</i>	<i>observations</i>	
	FA 6	FA 7
To a 1 cm depth of solution in a test-tube, add an equal volume of FA 3 , aqueous potassium iodide, followed by a few drops of starch indicator.	X	
To a 1 cm depth of solution in a boiling tube, add aqueous sodium hydroxide, then		
heat gently and carefully .		X
To a 1 cm depth of solution in a test-tube, add a few drops of aqueous silver nitrate.		
To a 1 cm depth of solution in a test-tube, add aqueous ammonia.		
To a 1 cm depth of solution in a test-tube, add a folded 3 cm length of magnesium ribbon.		

- (ii)** What can you deduce about solution **FA 7** from its reaction with magnesium?
Explain your answer.

.....

.....

- (iii) Give the ionic equation for the reaction of the metal cation in **FA 7** with aqueous sodium hydroxide. Include state symbols.

.....

- (iv) What **type** of reaction took place when aqueous potassium iodide was added to **FA 7**? Use your observations to help you explain your answer.

.....

.....

.....

- (v) The observation you made when aqueous silver nitrate was added to **FA 7** does not allow the anion in **FA 7** to be identified with certainty.

Explain why you cannot be certain about the identity of the anion.

.....

.....

.....

- (vi) A student suggested that the anion in **FA 7** could be identified with more certainty if excess ammonia solution was added after the aqueous silver nitrate.

Explain why this suggestion is **not** correct.

.....

.....

.....

[11]

[Total: 14]

Qualitative Analysis Notes

1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on heating	–
barium, Ba ²⁺ (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$)
bromide, $\text{Br}^-(\text{aq})$	gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$)
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$)
nitrate, $\text{NO}_3^-(\text{aq})$	NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and <i>Al</i> foil
nitrite, $\text{NO}_2^-(\text{aq})$	NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and <i>Al</i> foil; NO liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown NO_2 in air)
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)

3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint

The Periodic Table of Elements

		Group															
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">1 H hydrogen 1.0</div> <div style="border: 1px solid black; padding: 2px;">2 He helium 4.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">3 Li lithium 6.9</div> <div style="border: 1px solid black; padding: 2px;">4 Be beryllium 9.0</div> <div style="border: 1px solid black; padding: 2px;">5 B boron 10.8</div> <div style="border: 1px solid black; padding: 2px;">6 C carbon 12.0</div> <div style="border: 1px solid black; padding: 2px;">7 N nitrogen 14.0</div> <div style="border: 1px solid black; padding: 2px;">8 O oxygen 16.0</div> <div style="border: 1px solid black; padding: 2px;">9 F fluorine 19.0</div> <div style="border: 1px solid black; padding: 2px;">10 Ne neon 20.2</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">11 Na sodium 23.0</div> <div style="border: 1px solid black; padding: 2px;">12 Mg magnesium 24.3</div> <div style="border: 1px solid black; padding: 2px;">13 Al aluminium 27.0</div> <div style="border: 1px solid black; padding: 2px;">14 Si silicon 28.1</div> <div style="border: 1px solid black; padding: 2px;">15 P phosphorus 31.0</div> <div style="border: 1px solid black; padding: 2px;">16 S sulfur 32.1</div> <div style="border: 1px solid black; padding: 2px;">17 Cl chlorine 35.5</div> <div style="border: 1px solid black; padding: 2px;">18 Ar argon 39.9</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">19 K potassium 39.1</div> <div style="border: 1px solid black; padding: 2px;">20 Ca calcium 40.1</div> <div style="border: 1px solid black; padding: 2px;">21 Sc scandium 45.0</div> <div style="border: 1px solid black; padding: 2px;">22 Ti titanium 47.9</div> <div style="border: 1px solid black; padding: 2px;">23 V vanadium 50.9</div> <div style="border: 1px solid black; padding: 2px;">24 Cr chromium 52.0</div> <div style="border: 1px solid black; padding: 2px;">25 Mn manganese 54.9</div> <div style="border: 1px solid black; padding: 2px;">26 Fe iron 55.8</div> <div style="border: 1px solid black; padding: 2px;">27 Co cobalt 58.9</div> <div style="border: 1px solid black; padding: 2px;">28 Ni nickel 58.7</div> <div style="border: 1px solid black; padding: 2px;">29 Cu copper 63.5</div> <div style="border: 1px solid black; padding: 2px;">30 Zn zinc 65.4</div> <div style="border: 1px solid black; padding: 2px;">31 Ga gallium 69.7</div> <div style="border: 1px solid black; padding: 2px;">32 Ge germanium 72.6</div> <div style="border: 1px solid black; padding: 2px;">33 As arsenic 74.9</div> <div style="border: 1px solid black; padding: 2px;">34 Se selenium 79.0</div> <div style="border: 1px solid black; padding: 2px;">35 Br bromine 79.9</div> <div style="border: 1px solid black; padding: 2px;">36 Kr krypton 83.8</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">37 Rb rubidium 85.5</div> <div style="border: 1px solid black; padding: 2px;">38 Sr strontium 87.6</div> <div style="border: 1px solid black; padding: 2px;">39 Y yttrium 88.9</div> <div style="border: 1px solid black; padding: 2px;">40 Zr zirconium 91.2</div> <div style="border: 1px solid black; padding: 2px;">41 Nb niobium 92.9</div> <div style="border: 1px solid black; padding: 2px;">42 Mo molybdenum 95.9</div> <div style="border: 1px solid black; padding: 2px;">43 Tc technetium —</div> <div style="border: 1px solid black; padding: 2px;">44 Ru ruthenium 101.1</div> <div style="border: 1px solid black; padding: 2px;">45 Rh rhodium 106.4</div> <div style="border: 1px solid black; padding: 2px;">46 Pd palladium 106.4</div> <div style="border: 1px solid black; padding: 2px;">47 Ag silver 107.9</div> <div style="border: 1px solid black; padding: 2px;">48 Cd cadmium 112.4</div> <div style="border: 1px solid black; padding: 2px;">49 In indium 114.8</div> <div style="border: 1px solid black; padding: 2px;">50 Sn tin 118.7</div> <div style="border: 1px solid black; padding: 2px;">51 Sb antimony 121.8</div> <div style="border: 1px solid black; padding: 2px;">52 Te tellurium 127.6</div> <div style="border: 1px solid black; padding: 2px;">53 I iodine 126.9</div> <div style="border: 1px solid black; padding: 2px;">54 Xe xenon 131.3</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">55 Cs caesium 132.9</div> <div style="border: 1px solid black; padding: 2px;">56 Ba barium 137.3</div> <div style="border: 1px solid black; padding: 2px;">57–71 lanthanoids</div> <div style="border: 1px solid black; padding: 2px;">72 Hf hafnium 178.5</div> <div style="border: 1px solid black; padding: 2px;">73 Ta tantalum 180.9</div> <div style="border: 1px solid black; padding: 2px;">74 W tungsten 183.8</div> <div style="border: 1px solid black; padding: 2px;">75 Re rhenium 186.2</div> <div style="border: 1px solid black; padding: 2px;">76 Os osmium 190.2</div> <div style="border: 1px solid black; padding: 2px;">77 Ir iridium 192.2</div> <div style="border: 1px solid black; padding: 2px;">78 Pt platinum 195.1</div> <div style="border: 1px solid black; padding: 2px;">79 Au gold 197.0</div> <div style="border: 1px solid black; padding: 2px;">80 Hg mercury 200.6</div> <div style="border: 1px solid black; padding: 2px;">81 Tl thallium 204.4</div> <div style="border: 1px solid black; padding: 2px;">82 Pb lead 207.2</div> <div style="border: 1px solid black; padding: 2px;">83 Bi bismuth 209.0</div> <div style="border: 1px solid black; padding: 2px;">84 Po polonium —</div> <div style="border: 1px solid black; padding: 2px;">85 At astatine —</div> <div style="border: 1px solid black; padding: 2px;">86 Rn radon —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">87 Fr francium —</div> <div style="border: 1px solid black; padding: 2px;">88 Ra radium —</div> <div style="border: 1px solid black; padding: 2px;">89–103 actinoids</div> <div style="border: 1px solid black; padding: 2px;">104 Rf rutherfordium —</div> <div style="border: 1px solid black; padding: 2px;">105 Db dubnium —</div> <div style="border: 1px solid black; padding: 2px;">106 Sg seaborgium —</div> <div style="border: 1px solid black; padding: 2px;">107 Bh bohrium —</div> <div style="border: 1px solid black; padding: 2px;">108 Hs hassium —</div> <div style="border: 1px solid black; padding: 2px;">109 Mt meitnerium —</div> <div style="border: 1px solid black; padding: 2px;">110 Ds darmstadtium —</div> <div style="border: 1px solid black; padding: 2px;">111 Rg roentgenium —</div> <div style="border: 1px solid black; padding: 2px;">112 Cn copernicium —</div> <div style="border: 1px solid black; padding: 2px;">114 Fl flerovium —</div> <div style="border: 1px solid black; padding: 2px;">116 Lv livermorium —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">57 La lanthanum 138.9</div> <div style="border: 1px solid black; padding: 2px;">58 Ce cerium 140.1</div> <div style="border: 1px solid black; padding: 2px;">59 Pr praseodymium 140.9</div> <div style="border: 1px solid black; padding: 2px;">60 Nd neodymium 144.4</div> <div style="border: 1px solid black; padding: 2px;">61 Pm promethium —</div> <div style="border: 1px solid black; padding: 2px;">62 Sm samarium 150.4</div> <div style="border: 1px solid black; padding: 2px;">63 Eu europium 152.0</div> <div style="border: 1px solid black; padding: 2px;">64 Gd gadolinium 157.3</div> <div style="border: 1px solid black; padding: 2px;">65 Tb terbium 158.9</div> <div style="border: 1px solid black; padding: 2px;">66 Dy dysprosium 162.5</div> <div style="border: 1px solid black; padding: 2px;">67 Ho holmium 164.9</div> <div style="border: 1px solid black; padding: 2px;">68 Er erbium 167.3</div> <div style="border: 1px solid black; padding: 2px;">69 Tm thulium 168.9</div> <div style="border: 1px solid black; padding: 2px;">70 Yb ytterbium 173.1</div> <div style="border: 1px solid black; padding: 2px;">71 Lu lutetium 175.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">89 Ac actinium —</div> <div style="border: 1px solid black; padding: 2px;">90 Th thorium 232.0</div> <div style="border: 1px solid black; padding: 2px;">91 Pa protactinium 231.0</div> <div style="border: 1px solid black; padding: 2px;">92 U uranium 238.0</div> <div style="border: 1px solid black; padding: 2px;">93 Np neptunium —</div> <div style="border: 1px solid black; padding: 2px;">94 Pu plutonium —</div> <div style="border: 1px solid black; padding: 2px;">95 Am americium —</div> <div style="border: 1px solid black; padding: 2px;">96 Cm curium —</div> <div style="border: 1px solid black; padding: 2px;">97 Bk berkelium —</div> <div style="border: 1px solid black; padding: 2px;">98 Cf californium —</div> <div style="border: 1px solid black; padding: 2px;">99 Es einsteinium —</div> <div style="border: 1px solid black; padding: 2px;">100 Fm fermium —</div> <div style="border: 1px solid black; padding: 2px;">101 Md mendelevium —</div> <div style="border: 1px solid black; padding: 2px;">102 No nobelium —</div> <div style="border: 1px solid black; padding: 2px;">103 Lr lawrencium —</div> </div>															

lanthanoids

actinoids