
PHYSICS

9702/21

Paper 2 AS Level Structured Questions

May/June 2019

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **10** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Question	Answer	Marks
1(a)	(velocity =) change in displacement / time (taken)	B1
1(b)(i)	$k = [1.29 \times (3.3 \times 10^2)^2] / 9.9 \times 10^4$	C1
	= 1.4	A1
1(b)(ii)	percentage uncertainty = $(3 \times 2) + 4 + 2$ (= 12%) or fractional uncertainty = $(0.03 \times 2) + 0.04 + 0.02$ (= 0.12)	C1
	$\Delta k = 0.12 \times 1.42$ = 0.17 (<i>allow to 1 significant figure</i>)	C1
	$k = 1.4 \pm 0.2$	A1

PUBLISHED

Question	Answer	Marks
2(a)	(momentum =) mass \times velocity	B1
2(b)(i)	time = 40 ms	A1
2(b)(ii)	1. (the magnitude of the acceleration is) constant	B1
	2. (the magnitude of the acceleration is) zero	B1
2(c)	$F = \Delta p / (\Delta)t$ or $F = \text{gradient}$	C1
	e.g. $F = 0.50 / 40 \times 10^{-3}$ $= 13 \text{ N}$	A1
2(d)	horizontal line from (0, 0.40) to (60, 0.40)	B1
	straight line from (60, 0.40) to (100, -0.10)	B1
	horizontal line from (100, -0.10) to (160, -0.10)	B1

Question	Answer	Marks
3(a)(i)	$E = \frac{1}{2}Fx$ or $E = \frac{1}{2}kx^2$ or $E = \text{area under graph}$	C1
	$E = \frac{1}{2} \times 4.0 \times 0.32 = 0.64 \text{ J}$ or $E = \frac{1}{2} \times 12.5 \times (0.32)^2 = 0.64 \text{ J}$	A1
3(a)(ii)	$E = mgh$ or $E = Wh$ $= 2.5 \times 0.32$	C1
	$= 0.80 \text{ J}$	A1
3(b)(i)	kinetic energy $= 0.80 - 0.64$ $= 0.16 \text{ J}$	A1
3(b)(ii)	$E = \frac{1}{2}mv^2$ $0.16 = \frac{1}{2} \times (2.5 / 9.81) \times v^2$	C1
	$v = 1.1 \text{ m s}^{-1}$	A1

PUBLISHED

Question	Answer	Marks
4(a)(i)	$m = \rho \times V$	C1
	$= (4 / 3) \times \pi \times (1.2 \times 10^{-6})^3 \times 940 = 6.8 \times 10^{-15} \text{ (kg)}$	A1
4(a)(ii)	minimum charge (on drop) is $1.6 \times 10^{-19} \text{ C}$	B1
4(b)(i)	$V = Ed$	C1
	$V = 2.1 \times 10^5 \times 8.0 \times 10^{-3}$	A1
	$= 1.7 \times 10^3 \text{ V}$	
4(b)(ii)	constant velocity so no resultant force (so in equilibrium)	B1
4(b)(iii)	$mg = Eq$ or $mg = (V/d)q$ or $F = mg$ and $F = Eq$	C1
	$q = (6.8 \times 10^{-15} \times 9.81) / 2.1 \times 10^5$ $= 3.2 \times 10^{-19} \text{ C}$	A1
	sign of charge is negative	A1
4(c)(i)	<u>electric</u> force decreases	B1
	weight > electric force or resultant force acts downwards	B1
4(c)(ii)	(field line) separation increases	B1
4(d)(i)	upthrust (force)	B1
4(d)(ii)	air resistance/drag/viscous (force)	B1

Question	Answer	Marks
5(a)(i)	1. $N\lambda$	B1
	2. N/f	B1
5(a)(ii)	v (= distance / time) = $N\lambda / (N/f)$ so $v = f\lambda$	B1
5(b)	$T = 4.0 \times 0.20 = 0.80$ (ms) or 8.0×10^{-4} (s)	C1
	$f = 1 / 8.0 \times 10^{-4}$ $= 1300$ Hz	A1
5(c)(i)	constant phase difference (between the waves)	B1
5(c)(ii)	180°	A1
5(c)(iii)	path difference = 2λ or $S_1Y - S_2Y = 2\lambda$	C1
	distance = $7.40 + (0.85 \times 2)$ $= 9.1$ m	A1

PUBLISHED

Question	Answer	Marks
6(a)	energy is dissipated in the internal resistance	B1
6(b)	$E = V + Ir$	B1
6(c)(i)	(graph shows) maximum value of potential difference is 2.8 (V) or (graph shows) when current I (from battery) is zero, V is $2.8 (V) / E$	B1
6(c)(ii)	$r = (-)\text{gradient}$ or $r = (E - V) / I$ or substituted values from the graph for E , V and I	C1
	$r = 1.4 \Omega$	A1
6(d)(i)	$R = 2.1 / 0.50$ $= 4.2 \Omega$	A1
6(d)(ii)	number = $0.50 / 1.60 \times 10^{-19}$ $= 3.1 \times 10^{18}$	A1
6(d)(iii)	energy = EIt	C1
	or $P = EI$ and $P = W / t$	
	$(9.2 - 1.6) \times 10^3 = 2.8 \times 0.50 \times t$	
	$t = 5.4 \times 10^3 \text{ s}$	A1

PUBLISHED

Question	Answer	Marks
7(a)	nucleus is charged	B1
	the mass is <u>concentrated</u> in (very small) nucleus or the <u>majority</u> of the mass is in (very small) nucleus	B1
7(b)(i)	$-(1/3)e$	B1
7(b)(ii)	$2q - (1/3)e = e$ so $q = (2/3)e$	M1
	up / u (quarks) (<i>allow charm or top quarks</i>)	A1