

Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

BIOLOGY 0610/63

Paper 6 Alternative to Practical

May/June 2017

MARK SCHEME
Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE®, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

® IGCSE is a registered trademark.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



Mark schemes will use these abbreviations

• ; separates marking points

/ alternatives

I ignoreR reject

• A accept (for answers correctly cued by the question, or guidance for examiners)

AW alternative wording (where responses vary more than usual)

• AVP any valid point

• ecf credit a correct statement / calculation that follows a previous wrong response

ora or reverse argument

• () the word / phrase in brackets is not required, but sets the context

• <u>underline</u> actual word given must be used by candidate (grammatical variants excepted)

• max indicates the maximum number of marks that can be given

© UCLES 2017 Page 2 of 5

Question	Answer		Marks	Guidance
1(a)	table with two columns ;		3	
	column 1 heading 'vegetable extract', with unit as cm ³ ;	column 2 headed volume of iodine and		
	correct readings from Fig 1.2;			
1(b)	to allow iodine to change colour ;		1	A as an indicator
1(c)	volume of vegetable extract;		2	I amount
	volume / concentration of starch solution;			R volume iodine solution
	concentration of iodine;			
	temperature;			
	mixing time ;			
1(d)			4	improvement must relate to given error
	source of error ;;	improvement ;;		
	contamination	washing all apparatus / use new syringes		
	overshoot of end-point / adding too much iodine / many drops	add smaller quantities of iodine		
	determination of end-point	allow sufficient time for colour to change / use colorimeter / colour standard		A subjective colour change
	change in vitamin C with time	test same time after extraction for each		
	no repeats	repeat each concentration		
	AVP e.g. difficult reading scale coloured vegetable extracts	AVP e.g. use burette		

© UCLES 2017 Page 3 of 5

Question	Answer	Marks	Guidance
1(e)(i)	L : 25.00 ;	3	
	N : 62.5 ;		
	correct number of decimal places on both ;		
1(e)(ii)	axes labelled and units ;	4	
	even scale to fill more than half of the printed grid;		
	plot three / four points correctly;		ecf candidate result for 1(e)(i)
	line of best fit / trend line ;		
1(e)(iii)	mark volume of iodine used on (<i>y</i> -axis of) graph / extended horizontally and extended line vertically from plotted point to <i>x</i> -axis;	2	
	correct reading from graph on answer line;		
1(f)	range of temperatures ;	6	minimum of three
	values for temperatures stated ; time at each temperature ;		at least one above 50
	use of water-bath / named method;		
	description of extracting juice ;		
	detail of use of iodine drops / volume / addition of starch for end point ;;		
	at least two repeats;		
	(controlled variables) heating time / same type of vegetable / all samples from same vegetable;		
	relevant reference to safety;		

© UCLES 2017 Page 4 of 5

Question	Answer	Marks	Guidance
2(a)	any four from: drawing with clear outline; scaled to fit more than half the space; shape 5/6 sides for both; detail showing 3/4 layers with no shading and no cells;	4	
2(b)(i)	length of PQ = 80 mm;	3	± 1mm
	(x)64 ;;		(80) ÷ 1.25
2(b)(ii)	plane of section / AW ; magnification ; number of villi different ;	2	
2(c)(i)	A: 3 B: 9 C: 11	1	A 2 instead of 3 for A 3 correct answers = 1 mark
2(c)(ii)	30 °C; has highest rate of reaction / AW;	2	
2(c)(iii)	it is much higher / different than trial 1 and 3 / AW;	1	
2(c)(iv)	(IV) temperature ; (DV) rate of reaction ;	2	

© UCLES 2017 Page 5 of 5