



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



CHEMISTRY

0620/32

Paper 3 (Extended)

May/June 2013

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **14** printed pages and **2** blank pages.



1 Air is a mixture of gases. The main constituents are the elements oxygen and nitrogen.

(a) (i) Name another element in air.

..... [1]

(ii) Give the formula of a compound in unpolluted air.

..... [1]

(b) Common pollutants present in air are the oxides of nitrogen and sulfur dioxide.

(i) How are the oxides of nitrogen formed?

.....
.....
..... [2]


(ii) How is sulfur dioxide formed?

.....
.....
..... [2]

(iii) These oxides are largely responsible for acid rain.
State **two** harmful effects of acid rain.

.....
..... [2]

(c) The percentage of oxygen in air can be determined by the following experiment.



The gas syringe contains 50 cm³ of air. The large pile of copper is heated and the air is passed from one gas syringe to the other over the hot copper. The large pile of copper turns black. The gas is allowed to cool and its volume measured.

The small pile of copper is heated and the remaining gas passed over the hot copper. The copper does not turn black. The final volume of gas left in the apparatus is less than 50 cm³.

(i) Explain why the copper in the large pile turns black.

.....
 [2]

(ii) Why must the gas be allowed to cool before its volume is measured?

..... [1]

(iii) Explain why the copper in the small pile did not turn black.

..... [1]

(iv) What is the approximate volume of the gas left in the apparatus?

..... [1]

[Total: 13]

- 2 (a) The table below gives the number of protons, neutrons and electrons in atoms or ions. Complete the table. The first line is given as an example. You will need to use the Periodic Table.

particle	number of protons	number of electrons	number of neutrons	symbol or formula
A	4	4	5	${}^9_4\text{Be}$
B	19	18	20
C	30	30	35
D	8	10	8
E	31	31	39

[6]


- (b) Using the data in the table, explain how you can determine whether a particle is an atom, a negative ion or a positive ion.

.....

 [3]

[Total: 9]

- 3 The diagram shows some of the processes which determine the percentage of carbon dioxide in the atmosphere.



- (a) Explain how the following two processes alter the percentage of carbon dioxide in the atmosphere.

- (i) combustion

.....

 [3]

(ii) respiration

.....
.....
..... [3]

(b) Photosynthesis reduces the percentage of carbon dioxide in the atmosphere.

(i) Complete the word equation for photosynthesis.

carbon dioxide + water → + [2]


(ii) State **two** essential conditions for the above reaction to occur.

.....
..... [2]


[Total: 10]

4 At present the most important method of manufacturing hydrogen is steam reforming of methane.

(a) In the first stage of the process, methane reacts with steam at 800 °C.



In the second stage of the process, carbon monoxide reacts with steam at 200 °C.



(i) Explain why the position of equilibrium in the first reaction is affected by pressure but the position of equilibrium in the second reaction is not.

.....
.....
..... [2]

(ii) Suggest why a high temperature is needed in the first reaction to get a high yield of products but in the second reaction a high yield is obtained at a low temperature.

.....
..... [2]

(b) Two other ways of producing hydrogen are cracking and electrolysis.

- (i) Hydrogen can be a product of the cracking of long chain alkanes. Complete the equation for the cracking of C_8H_{18} .



- (ii) There are three products of the electrolysis of concentrated aqueous sodium chloride. Hydrogen is one of them. Write an equation for the electrode reaction which forms hydrogen.

..... [2]

- (iii) Name the other **two** products of the electrolysis of concentrated aqueous sodium chloride and give a use of each one.


product use

product use [4]

[Total: 11]

5 Many monomer molecules react together to form one molecule of a polymer. This reaction is called polymerisation.

- (a) The structural formula of the polymer, poly(chloroethene), is given below. This polymer is also known as PVC.



- (i) A major use of PVC is insulation of electric cables. PVC is a poor conductor of electricity.

Suggest another property which makes it suitable for this use.


..... [1]

- (ii) One way of disposing of waste PVC is by burning it. This method has the disadvantage that poisonous gases are formed.

Suggest **two** poisonous gases which could be formed by the combustion of PVC.

..... [2]


- (b) (i) Deduce the structural formula of the monomer from that of the polymer.



structural formula of monomer

[1]

- (ii) Deduce the structural formula of the polymer, poly(phenylethene), from the formula of its monomer, phenylethene.




structural formula of polymer

[2]

- (c) The carbohydrate, glucose, polymerises to form the more complex carbohydrate starch.

If glucose is represented by



then the structural formula of starch is as drawn below.



How does the polymerisation of glucose differ from that of an alkene such as phenylethene?


.....

 [2]

[Total: 8]

- 6 Aluminium is an important metal with a wide range of uses.

- (a) Aluminium is obtained by the electrolysis of aluminium oxide dissolved in molten cryolite.



- (i) Solid aluminium oxide is a poor conductor of electricity. It conducts either when molten or when dissolved in molten cryolite. Explain why.

.....

 [2]

- (ii) Why is a solution of aluminium oxide in molten cryolite used rather than molten aluminium oxide?

..... [1]

(iii) Explain why the carbon anodes need to be replaced periodically.

..... [1]


(iv) One reason why graphite is used for the electrodes is that it is a good conductor of electricity. Give another reason.

..... [1]

(b) Aluminium is used to make food containers because it resists corrosion. Explain why it is not attacked by the acids in food.

..... [2]

(c) Aluminium is used for overhead power (electricity) cables which usually have a steel core.



(i) Give **two** properties of aluminium which make it suitable for this use.


..... [2]

(ii) Explain why the cables have a steel core.

..... [1]

[Total: 10]

7 The ester linkage showing all the bonds is drawn as



or more simply it can be written as -COO- .

(a) (i) Give the structural formula of the ester ethyl ethanoate.

[1]


(ii) Deduce the name of the ester formed from methanoic acid and butanol.

..... [1]

(b) (i) Which group of naturally occurring compounds contains the ester linkage?

..... [1]


(ii) Draw the structural formula of the polyester formed from the following monomers.



You are advised to use the simpler form of the ester linkage.

[3]

- (c) Esters can be used as solvents in chromatography. The following shows a chromatogram of plant acids.



An ester was used as the solvent and the chromatogram was sprayed with bromothymol blue.

- (i) Suggest why it was necessary to spray the chromatogram.

.....
 [2]

- (ii) Explain what is meant by the R_f value of a sample.

.....
 [1]

(iii) Calculate the R_f values of the two samples and use the data in the table to identify the plant acids.

plant acid	R_f value
tartaric acid	0.22
citric acid	0.30
oxalic acid	0.36
malic acid	0.46
succinic acid	0.60

sample 1 $R_f = \dots\dots\dots$ It is $\dots\dots\dots$ acid.

sample 2 $R_f = \dots\dots\dots$ It is $\dots\dots\dots$ acid. [2]

[Total: 11]

8 (a) Define the following

(i) the mole

.....
 [1]

(ii) the Avogadro constant

.....
 [1]


(b) Which **two** of the following contain the same number of molecules?
 Show how you arrived at your answer.

- 2.0 g of methane, CH₄
- 8.0 g of oxygen, O₂
- 2.0 g of ozone, O₃
- 8.0 g of sulfur dioxide, SO₂

.....

 [2]

(c) 4.8 g of calcium is added to 3.6 g of water. The following reaction occurs.



(i) the number of moles of Ca =

the number of moles of H₂O = [1]

(ii) Which reagent is in excess? Explain your choice.

.....
..... [2]

(iii) Calculate the mass of the reagent named in (ii) which remained at the end of the experiment.

.....
..... [1]

[Total: 8]

DATA SHEET
The Periodic Table of the Elements

Group		I	II	III	IV	V	VI	VII	0
		1 H Hydrogen 1							4 He Helium 2
7 Li Lithium 3	9 Be Beryllium 4	11 B Boron 5	12 C Carbon 6	13 Al Aluminium 13	14 N Nitrogen 7	15 O Oxygen 8	16 F Fluorine 9	17 Ne Neon 10	
23 Na Sodium 11	24 Mg Magnesium 12	27 Al Aluminium 13	28 Si Silicon 14	29 P Phosphorus 15	30 S Sulfur 16	31 Cl Chlorine 17	32 Ar Argon 18		
39 K Potassium 19	40 Ca Calcium 20	45 Sc Scandium 21	46 Ti Titanium 22	47 V Vanadium 23	48 Cr Chromium 24	49 Mn Manganese 25	50 Fe Iron 26	51 Co Cobalt 27	52 Ni Nickel 28
85 Rb Rubidium 37	86 Sr Strontium 38	87 Y Yttrium 39	88 Zr Zirconium 40	89 Nb Niobium 41	90 Mo Molybdenum 42	91 Tc Technetium 43	92 Ru Ruthenium 44	93 Rh Rhodium 45	94 Pd Palladium 46
133 Cs Caesium 55	137 Ba Barium 56	138 La Lanthanum 57	139 Ce Cerium 58	140 Pr Praseodymium 59	141 Nd Neodymium 60	142 Pm Promethium 61	143 Sm Samarium 62	144 Eu Europium 63	145 Gd Gadolinium 64
226 Fr Francium 87	227 Ra Radium 88	228 Ac Actinium 89	229 Th Thorium 90	230 Pa Protactinium 91	231 U Uranium 92	232 Np Neptunium 93	233 Pu Plutonium 94	234 Am Americium 95	235 Cm Curium 96
		59 Co Cobalt 27	60 Ni Nickel 28	61 Cu Copper 29	62 Zn Zinc 30	63 Ga Gallium 31	64 Ge Germanium 32	65 As Arsenic 33	66 Se Selenium 34
		71 In Indium 49	72 Sb Antimony 51	73 Te Tellurium 52	74 Pb Lead 82	75 Bi Bismuth 83	76 Po Polonium 84	77 At Astatine 85	78 Rn Radon 86
		101 Ru Ruthenium 44	102 Rh Rhodium 45	103 Pd Palladium 46	104 Ag Silver 47	105 Cd Cadmium 48	106 In Indium 49	107 Sn Tin 50	108 Pb Lead 82
		159 Tb Terbium 65	160 Dy Dysprosium 66	161 Ho Holmium 67	162 Er Erbium 68	163 Tm Thulium 69	164 Yb Ytterbium 70	165 Lu Lutetium 71	
		181 Ta Tantalum 73	182 Hf Hafnium 72	183 W Tungsten 74	184 Re Rhenium 75	185 Os Osmium 76	186 Ir Iridium 77	187 Pt Platinum 78	188 Au Gold 79
		201 Hg Mercury 80	202 Tl Thallium 81	203 Pb Lead 82	204 Bi Bismuth 83	205 Po Polonium 84	206 At Astatine 85	207 Rn Radon 86	
		269 Uu Ununennium 111	270 Uub Unbium 112	271 Uut Ununtrium 113	272 Uuq Unquadium 114	273 Uup Unpentium 115	274 Uuq Unseptium 116	275 Uuh Unhassium 117	276 Uu Unnium 118

***58-71 Lanthanoid series**
†90-103 Actinoid series

a	X
b	†

Key
a = relative atomic mass
X = atomic symbol
b = proton (atomic) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.