

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER		CANDIDATE NUMBER
CHEMISTRY		0620/6
Paper 6 Alterna	tive to Practical	May/June 201
		1 hou
Candidates ans	swer on the Question Paper.	
No Additional M	laterials are required.	

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **11** printed pages and **1** blank page.




1 Electricity was passed through a solution of concentrated hydrochloric acid using the apparatus shown.

For Examiner's Use



- 2 A student found a recipe for making elderberry wine by fermentation.
 - 1 kg elderberries 0.5 kg sugar 10 g yeast granules 3 dm³ water

The student decided to make some elderberry wine using the apparatus below.



The student carried out the following method.

- Step 1 The elderberries were crushed.
- Step 2 The crushed elderberries and sugar were added to the water and the mixture was boiled for ten minutes. The crushed elderberries were then separated from the mixture.
- Step 3 Yeast was added to the liquid when it had cooled to room temperature.
- (a) Suggest the purpose of the airlock in the apparatus.
-[1]
- (b) What apparatus could be used in Step 1?
 -[1]
- (c) Draw a labelled diagram of the apparatus used to separate the crushed elderberries from the mixture in Step 2.

[2]

(d) Why was the yeast in Step 3 not added until the liquid was at room temperature?

 [1]

 [1]

(e) (i) State one observation during the fermentation. Examiner's (ii) Suggest how the rate of the fermentation reaction could be measured. (f) Name the method that could be used to separate ethanol from the fermented mixture.[1] [Total: 9]

For

Use

4

3 A student investigated the reaction between two different solids, **C** and **D**, and excess dilute hydrochloric acid.

Five experiments were carried out.

(a) Experiment 1

A measuring cylinder was used to pour 30 cm^3 of dilute hydrochloric acid into a polystyrene cup. The temperature of the dilute hydrochloric acid was measured. 1 g of solid **C** was added to the dilute hydrochloric acid and the mixture stirred with a thermometer. The maximum temperature reached by the liquid mixture was measured.

(b) Experiment 2

The polystyrene cup was emptied and rinsed with water. Experiment 1 was repeated using 2 g of solid **C**.

(c) Experiments 3 and 4

Experiment 2 was repeated using 3 g and then 5 g of solid **C**.

Use the thermometer diagrams to record the results in the table below.

Complete the final column in the table.

experiment	mass of solid C /g	thermometer diagram	initial temperature of acid/°C	thermometer diagram	maximum temperature reached/°C	temperature difference /°C
1		30 25 20		30 -25 -20		
2		30 -25 -20		35 -30 -25		
3		30 -25 -20		35 -30 -25		
4		30 -25 -20		-35 -30 -25		

For

Examiner's

Use


[3]

(d) Experiment 5

Experiment 1 was repeated using solid **D**. Use the thermometer diagrams to record the results in the spaces below.

For

Examiner's Use



(f)	(i)	From your graph , deduce the temperature of the solution when 6g of solid C is added to 30 cm ³ of dilute hydrochloric acid. Show clearly on the grid how you worked out your answer.	For Examiner's Use
		°C [2]	
	(ii)	From your graph , deduce the mass of solid C that would give a temperature rise of $9 ^{\circ}$ C when added to 30cm^3 of dilute hydrochloric acid.	
(g)	Wh	at type of chemical process occurs when solid D reacts with dilute hydrochloric acid?	
(h)		ggest the effect on the results if Experiment 3 was repeated using 60 cm ³ of dilute rochloric acid.	
(i)	Pre	dict the temperature of the solution in Experiment 4 after 1 hour. Explain your answer.	
		101	
(j)		en carrying out the experiments, what would be one advantage and one disadvantage aking the temperature readings after exactly one minute?	
	adv	antage	
	disa	advantage	
		[Total: 20]	


A mixture of two solids, E and F, was analysed.
 Solid E was the water-soluble salt aluminium chloride, AlCl₃, and solid F was an insoluble salt.

The tests on the mixture and some of the observations are in the following table. Complete the observations in the table.

	tests	observations
boiling The c and fil	ed water was added to the mixture in a g tube. ontents of the boiling tube were shaken Itered, keeping the filtrate and residue for llowing tests.	
tests o	on the filtrate	
	iltrate was divided into five portions in est-tubes.	
	he first portion was used to describe the ppearance of the filtrate.	appearance [1]
h p E	everal drops of aqueous sodium ydroxide were added to the second ortion of the solution. xcess aqueous sodium hydroxide was hen added to the test-tube.	[3]
1	queous ammonia was added to the third ortion, dropwise and then in excess.	
n	o the fourth portion of the solution, dilute itric acid and aqueous silver nitrate were dded.	
1	o the fifth portion of the solution, about cm ³ of dilute nitric acid and barium itrate solution were added.	[1]

tests	observations	
tests on the residue		
(f) (i) To a little of the residue, dilute	rapid effervescence	
hydrochloric acid was added. The gas given off was tested.	gas turned limewater milky	
(ii) The residue was heated, gently then strongly.	solid changed colour from green to black	
(g) What conclusions can you draw about s	solid F?	
	[Total: 11]	

5 Identical pieces of steel were placed in two different boiling liquids for 12 hours. The graphs show how the mass of each piece of steel changed.



10

For

Examiner's

Use

Copper(II) oxide and carbon are both black solids. Copper(II) oxide reacts with dilute sulfuric 6 Examiner's acid to form aqueous copper(II) sulfate. Carbon does not react with dilute sulfuric acid. You are given a mixture of copper(II) oxide and carbon and access to dilute sulfuric acid. Plan an experiment to investigate the percentage of copper(II) oxide in the mixture.

.....[6]

[Total: 6]

For

Use

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.