

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the October/November 2014 series

0620 CHEMISTRY

0620/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

bestexamhelp.com

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0620	31

1 (a) Match the following pH values to the solutions given below.

1 3 7 10 13

The solutions all have the same concentration.

solution	pH	
aqueous ammonia, weak base	10	
dilute hydrochloric acid, a strong acid	1	
aqueous sodium hydroxide, a strong base	13	
aqueous sodium chloride, a salt	7	
dilute ethanoic acid, a weak acid	3	[5]

(b) Hydrochloric acid strong acid **or** ethanoic acid weak acid [1]
OR: hydrochloric acid completely ionised **or** ethanoic acid partially ionised
hydrochloric acid greater concentration of/more H⁺ ions (than ethanoic acid) [1]

(c) Rate of reaction with Ca, Mg, Zn, Fe [1]

Strong (hydrochloric) acid bubbles faster **or** more bubbles **or** dissolves faster [1]

OR: rate of reaction with (metal) carbonate [1]

strong (hydrochloric) acid faster **or** more bubbles **or** dissolves faster (only if carbonate insoluble) [1]

OR: electrical conductivity [1]

strong (hydrochloric) acid better conductor [1]

[Total: 9]

2 (a) soft because weak forces between layers/sheets/rows [1]

layers can slip/slide [1]

good conductor because electrons can move/mobile [1]

(b) it is soft: pencils **or** lubricant **or** polish [1]

good conductor: electrodes **or** brushes (in electric motors) [1]

(c) (i) every silicon atom is bonded/attached to 4 oxygen atoms or every oxygen bonded/attached to two silicon atoms [1]

(ii) Any **two** from:
high melting point/boiling point
hard
colourless crystals/shiny
poor/non-conductor of electricity/insulator
insoluble in water [2]

[Total: 8]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0620	31

- 3 (a) Any **two** from:
bleach/making wood pulp/making paper
food/fruit juice/wine preservative
fumigant/sterilising/insecticide [2]
- (b) heating/roasting/burning (zinc sulfides) [1]
in air/oxygen COND on M1 [1]
- (c) (i) V_2O_5 [1]
- (ii) position of equilibrium shifts right/yield increases [1]
to save energy [1]
- (iii) faster reaction/rate [1]
more collisions per second/higher collision frequency [1]
fewer moles/molecules (of gas) on right [1]
(so) position of equilibrium shifts right/yield increases [1]
- (d) (the reaction is) too violent/too exothermic **or** produces mist/fumes (of acid) [1]
- [Total: 12]
- 4 (a) (i) insufficient/limited oxygen [1]
or $2C + O_2 \rightarrow 2CO$
- coke/carbon reacts with carbon dioxide [1]
or $C + CO_2 \rightarrow 2CO$
- (ii) $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$
species (1) balancing (1) [2]
- (b) (i) carbon dioxide [1]
- (ii) $CaO + SiO_2 \rightarrow CaSiO_3$ [2]
[1] each side correct
- (iii) (molten) iron higher density (than slag) [2]
- (iv) No oxygen in contact with iron **or** layer of slag prevents hot iron reacting with oxygen/air **or** (all) oxygen reacts with carbon (so no oxygen left to react with iron) [1]
- (c) (i) air/oxygen and water (need both) [1]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0620	31

(ii) aluminium oxide layer is impervious **or** non-porous **or** passive **or** unreactive **or** will not allow water/air to pass through it (rust allows passage of water **or** air **or** it flakes off) [1]

(d) (i) zinc more reactive (than iron/steel) [1]
loses electrons [1]
electrons move (from zinc) to iron [1]
Zinc reacts (with air and water) **or** zinc corrodes **or** zinc is oxidised **or** zinc is anodic **or** zinc forms positive ions **or** zinc forms Zn^{2+} **or** iron and steel don't react with air/water **or** iron and steel are not oxidised **or** iron and steel do not form ions **or** iron and steel do not lose electrons **or** iron and steel are cathodic [1]

(ii) R to L in wire [1]

(iii) $2H^+ + 2e^- \rightarrow H_2$
species (1) balancing (1)

[Total: 19]

5 (a) nitrogen and oxygen react [1]
at high temperatures (in engine) [1]

(b) M1 carbon monoxide (converted to) carbon dioxide **or** $2CO + O_2 \rightarrow 2CO_2$ [1]

M2 (by) oxides of nitrogen (which are reduced to) nitrogen
or $2NO \rightarrow N_2 + O_2$ **or** $2NO_2 \rightarrow N_2 + 2O_2$ [1]

M3 hydrocarbons (burn) making water [1]

M4 products: any **two** from:
carbon dioxide, water, nitrogen [1]

(c) lead compounds are toxic **or** brain damage **or** reduce IQ or nausea or kidney failure **or** anaemia [1]

[Total: 7]

6 (a) (i) butanoic acid [1]
methanol [1]

(ii) number of moles of ethanoic acid = 0.1 [1]
number of moles of ethanol = 0.12(0) [1]
the limiting reagent is ethanoic acid [1]
number of moles of ethyl ethanoate formed = 0.1 [1]
maximum yield of ethyl ethanoate is 8.8 g [1]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0620	31

(b) correct ester linkage [1]
 two ester linkages (COND on M1) [1]
 continuation (COND on M2) [1]

(c) (i) add bromine water/bromine [1]
 turns colourless [1]
 remains brown/orange/reddish brown/yellow [1]

ALLOW: potassium manganate(VII) (acidic or alkaline) [1]
 correct colour colourless/green or brown ppt [1]
 stays pink/purple [1]

(ii) ester 1 [1]
COND alkyl group is C_nH_{2n+1} which is NOT $C_{17}H_{33}$ [1]
or $C_{17}H_{35}$ is C_nH_{2n+1} **or** less hydrogen [1]

(iii) soap **or** (sodium) salt (of a carboxylic acid) **or** carboxylate [1]
 alcohol [1]

[Total: 17]

7 (a) (i) $6Li + N_2 = 2Li_3N$
 species (1) balancing (1)

(ii) N^{3-} ion drawn correctly [1]
 Charges correct (minimum 1 × Li ion and 1 nitride ion) [1]

(b) (i) 3 × shared pairs between N and 3 × F [1]
 only 2 non-bonding electrons on N, 6 non-bonding electrons on each F
 (COND on first point) [1]

(ii) Strong attractive forces/strong ionic bonds in lithium nitride [1]
 weak (attractive) forces between molecules in NF_3 [1]

[Total: 8]