

## Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

|         | CANDIDATE<br>NAME             |   |                          |
|---------|-------------------------------|---|--------------------------|
|         | CENTRE<br>NUMBER              | CANDIDATE<br>NUMBER                         |                          |
| * 0 8 5 | CHEMISTRY<br>Paper 6 Alternat | ive to Practical                            | 0620/62<br>May/June 2015 |
| 8696    | ·                             | wer on the Question Paper.                  | 1 hour                   |
| 3 1 5 * |                               | aterials are required.<br>MODIFIED LANGUAGE |                          |
|         |                               |   |                          |

## READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.Write in dark blue or black pen.You may use an HB pencil for any diagrams or graphs.Do not use staples, paper clips, glue or correction fluid.DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 11 printed pages and 1 blank page.



1 A student did an experiment to measure the energy produced by burning ethanol. The apparatus used is shown.



The ethanol was burned for one minute. The temperature of the water was then measured and recorded.

(a) Complete the boxes to name the pieces of apparatus.

[2]

(b) Give three other measurements the student should have taken.

| 1 | <br> | <br> | <br> |     |
|---|------|------|------|-----|
| 2 | <br> | <br> | <br> |     |
| 3 | <br> | <br> | <br> |     |
|   |      |      |      | [3] |

(c) The experiment was repeated using 40 cm<sup>3</sup> of water. What effect would this have on the results?

......[1]

(d) Another student did this experiment using a copper can instead of a boiling tube. Give one advantage of this change to the apparatus.

......[1]

[Total: 7]

**2** A student prepared some crystals of chromium(III) nitrate,  $Cr(NO_3)_3.6H_2O$ . The following extract was taken from his practical notes.

| _   |       |   |
|-----|-------|---|
|     |       | Making chromium(III) nitrate crystals   |
|     | Step  | 1 I poured 50 cm <sup>3</sup> of acid into a beaker. Solid chromium(III) oxide was then added a little at a time and the mixture stirred. |
|     | Step  | 2 When no more chromium(III) oxide reacted I separated the mixture and collected the solution in an evaporating dish.                     |
|     | Step  | 3 I boiled the solution strongly for ten minutes.   |
| (a) | Nam   | ne the acid used in this preparation.   |
|     |       |   |
| (b) | \\/bc | at would be used in Step 1 to   |
| (0) | VVIIC |   |
|     | (i)   | add the chromium(III) oxide to the acid,  |
|     |       |   |
|     | (ii)  | stir the mixture?   |
|     |       |   |
|     |       |   |
| (c) | Nam   | ne the separation method used in Step 2.  |
|     |       | [1]   |
|     |       |   |
| (d) | (i)   | Suggest what was left in the evaporating dish at the end of Step 3.   |
|     |       |   |
|     |       | How should the student have changed the method in Step 3 to obtain pure, dry crystals of chromium(III) nitrate?                           |
|     |       |   |
|     |       |   |
|     |       |   |
|     |       | [3]   |
|     |       | [Total: 8]  |
|     |       |   |

- 3 Three bottles of liquid have lost their labels. The liquids are known to be:
  - aqueous potassium hydroxide,
  - octane,
  - pure water.

Outline tests you would do to identify and distinguish the liquid in each bottle.


| liquid                      | test | result |
|-----------------------------|------|--------|
| aqueous potassium hydroxide |      |        |
| octane                      |      |        |
| pure water                  |      |        |

[6]

[Total: 6]

4 A student investigated the rate of reaction between hydrochloric acid and aqueous sodium thiosulfate. When these chemicals react they form a precipitate which makes the solution go cloudy. The formation of this precipitate can be used to show how fast the reaction proceeds.

Five experiments were done using the apparatus shown below.



(a) Experiment 1

Using a measuring cylinder, 50 cm<sup>3</sup> of aqueous sodium thiosulfate was poured into a conical flask. The conical flask was placed on a printed sheet of paper. 10 cm<sup>3</sup> of the hydrochloric acid was added to the solution in the conical flask and the stop clock started.

The time taken for the printed words to disappear from view was measured.

(b) Experiment 2

Using a measuring cylinder, 40 cm<sup>3</sup> of aqueous sodium thiosulfate was poured into a conical flask, followed by 10 cm<sup>3</sup> of distilled water. The conical flask was placed on the printed sheet. 10 cm<sup>3</sup> of the hydrochloric acid was added to the solution in the conical flask and the stop clock started.

The time taken for the printed words to disappear from view was measured.

(c) Experiment 3

Experiment 2 was repeated using 35 cm<sup>3</sup> of aqueous sodium thiosulfate and 15 cm<sup>3</sup> of distilled water.

(d) Experiment 4

Experiment 2 was repeated using 30 cm<sup>3</sup> of aqueous sodium thiosulfate and 20 cm<sup>3</sup> of distilled water.

(e) Experiment 5

Experiment 2 was repeated using 20 cm<sup>3</sup> of aqueous sodium thiosulfate and 30 cm<sup>3</sup> of distilled water.


(f) Use the stop clock diagrams to record the times in the table. Complete the table.

| Experiment<br>number | volume of<br>aqueous sodium<br>thiosulfate/cm <sup>3</sup> | volume of<br>distilled<br>water/cm <sup>3</sup> | stop clock<br>diagram  | time for<br>printed words<br>to disappear/s |
|----------------------|--|---|--|---|
| 1                    |  |   | 45<br>45<br>45<br>45<br>45<br>45<br>45<br>5<br>15<br>minutes |   |
| 2                    |  |   |  |   |
| 3                    |  |   |  |   |
| 4                    |  |   |  |   |
| 5                    |  |   |  |   |

[4]

7

(g) Plot the results on the grid and draw a smooth line graph.



- [3]
- (h) (i) From your graph, deduce the time for the printed words to disappear if Experiment 2 was repeated using 25 cm<sup>3</sup> of aqueous sodium thiosulfate and 25 cm<sup>3</sup> of distilled water. Show clearly on the grid how you worked out your answer.

......[3]

(ii) Sketch on the grid the curve you would expect if the experiments were repeated at a lower temperature. Label this curve 'lower temperature'. [1]

| (i) | (i)  | In which experiment was the rate of reaction greatest?                               |
|-----|------|--|
|     | (ii) | Explain why the rate of reaction was greatest in this experiment.                    |
|     |      | [1]  |
| (j) | A st | udent did a sixth experiment using 60 cm <sup>3</sup> of aqueous sodium thiosulfate. |
|     | Wh   | y would this not be an appropriate volume to use in this series of experiments?      |
|     |      |  |
| (k) | Sug  | gest and explain the effect of   |
|     | (i)  | using a burette to measure the volume of the hydrochloric acid,                      |
|     |      | [2]  |
|     | (ii) | using a 100 cm <sup>3</sup> conical flask.   |
|     |      |  |
|     |      |  |
|     |      | [Total: 19]  |


8

© UCLES 2015

Complete the observations in the table.

|                          | tests  | observations                                  |
|--------------------------|--|---|
| (a) Appea                | rance of the mixture.  | white solid                                   |
|                          | ter was added to the mixture.<br>e was shaken and filtered.  |   |
| tests on the             | e filtrate   |   |
|                          | on was divided into two<br>ons in two test-tubes.  |   |
| aqueo<br>added<br>gently | first portion of the solution,<br>us sodium hydroxide was<br>. The mixture was heated<br>and the gas evolved was<br>with pH indicator paper. | [2]   |
| solutio                  | second portion of the<br>n, dilute nitric acid and<br>us silver nitrate solution<br>added.   | [2]   |
| tests on the             | <u>e residue</u>   |   |
| to the                   | hydrochloric acid was added<br>residue in a test-tube. The<br>ven off was tested.  | rapid effervescence<br>limewater turned milky |
|                          | sulfuric acid was added to lution formed.  | white precipitate formed                      |
| <b>(e)</b> Wł            | nat is the pH value of the gas g   | given off in test <b>(b)</b> ?                |
|                          |  | [1]   |
| <b>(f)</b> Ide           | entify the gas given off in test (   |   |
| <br>(g) Wł               | nat are your conclusions about   | [1]<br>solid <b>K</b> ?                       |
|                          |  | [2]   |
|                          |  | [Total: 8]                                    |

6 Concentrated hydrochloric acid was electrolysed.



Hydrogen gas formed at the cathode (negative electrode).

| <b>(a)</b> Na | me a suitable metal to use for the electrodes.  |
|---------------|---|
|               |   |
| <b>(b)</b> Wh | y does hydrogen form at the negative electrode?   |
|               |   |
| (c) (i)       | Identify the gas given off at the anode (positive electrode).                                   |
|               | [1]   |
| (ii)          | Give a test for this gas.   |
|               | test  |
|               | result[2]   |
|               |   |
|               | ggest why the volume of gas formed at the positive electrode is less than the volume of drogen. |
|               |   |
|               | [Total: 6]  |
|               |   |

## **Tonic Water**

Tonic water is a solution containing citric acid. The concentration of the acid can be determined by reaction with aqueous potassium hydroxide solution.

Plan an investigation to show which of two different brands of colourless tonic water, Tastyton and Slimton, contains the highest concentration of citric acid.

You can use common laboratory apparatus and chemicals.

[6]

0620/62/M/J/15

7

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.