

## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/62
Paper 6 Alterna	ative to Practical		May/June 2016
			1 hour
Candidates ans	swer on the Question Paper.		
No Additional M	Naterials are required		

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1 The diagram shows the apparatus used to reduce copper(II) oxide with hydrogen.



(a)	Cor	nplete the boxes to name the apparatus.	[2]
(b)	Use	an arrow to indicate where heat is applied.	[1]
(c)	The	colour of the copper(II) oxide changes from to	[2]
(d)	Sug	gest a reason why the U-tube is surrounded by ice.	
			[1]
(e)	(i)	Identify the colourless liquid formed.	
			[1]
	(ii)	Give a chemical test for this liquid.	
		test	
		result	
			[2]
(	(iii)	How could you show that this liquid is pure?	
			[1]

[Total: 10]

- 2 A student investigated the rate of reaction between hydrogen peroxide and aqueous potassium iodide. When these chemicals react they form iodine. Sodium thiosulfate solution reacts with iodine and can be used to show how fast the reaction proceeds.
  - (a) A burette was filled up to the 0.0 cm<sup>3</sup> mark with sodium thiosulfate solution.

Using a large measuring cylinder, 100 cm³ of distilled water were poured into a conical flask. Using a small measuring cylinder, 6 cm³ of sulfuric acid, 1 cm³ of starch solution and 4 cm³ of aqueous potassium iodide were added to the flask.

0.5 cm<sup>3</sup> of sodium thiosulfate solution was added from the burette to the mixture in the flask and swirled to mix.

The reaction was then started by adding 3 cm<sup>3</sup> of hydrogen peroxide solution to the mixture, and the timer started.

The time taken for a blue colour to appear was noted.


A further  $0.5 \, \mathrm{cm^3}$  of sodium thiosulfate solution was added to the mixture in the conical flask, swirled and the blue colour disappeared. The time when the blue colour reappeared was noted. The experiment continued by adding further  $0.5 \, \mathrm{cm^3}$  portions of sodium thiosulfate solution until a total of  $3.0 \, \mathrm{cm^3}$  of sodium thiosulfate solution had been added, noting the times at which the blue colour reappeared.

Use the timer diagrams on page 4 to record the times in seconds in the table.

total volume of sodium thiosulfate solution added/cm <sup>3</sup>	timer diagram	time at which blue colour appeared/s
0.5	0 seconds 0 5 15 15 minutes	
1.0	45 15 15 15	
1.5	45 15 15	
2.0	45 15 15 15	
2.5	45 5 15	
3.0	45 15 15	

[3]

(b) Plot the results you have obtained on the grid and draw a best-fit straight-line graph.



time/s

volume of sodium thiosulfate solution/cm3

[5]

(c) (i) From your graph deduce the time at which the blue colour would appear if a total of 4.0 cm³ of sodium thiosulfate solution were added to the mixture in the conical flask. Show clearly on the grid how you worked out your answer.

[3]

(ii) Sketch on the grid the graph you would expect if the experiment was repeated at a higher temperature.
[1]

(d)	Su	ggest the purpose of the starch solution.
		[1]
(e)	(i)	Suggest <b>one</b> advantage of using a pipette to measure the volume of the hydrogen peroxide.
		[1]
	(ii)	Suggest and explain <b>one</b> disadvantage of using a pipette to measure the volume of the hydrogen peroxide.
		[2]
(f)	Exp	plain one disadvantage of using a beaker instead of a conical flask.
	••••	
		[1]
		[Total: 17]

3	Two solids, <b>E</b> and <b>F</b> , were analysed. Solid <b>E</b> was sodium sulfite. Both solids were found to be water
	soluble.

The tests on the solids, and some of the observations, are shown below.

4 4 -		1	12.0	
tests	_n	20	IIA.	_
ıcata	VII	30	IIU	_

tes	ts o	n solid E	
(a)	Des	scribe the appearance of the solid.	
			[1]
(b)	Dis	tilled water was added to solid <b>E</b> in a tes	st-tube and shaken to dissolve.
	The	e solution was divided into two portions in	n two test-tubes and the following tests carried out.
	(i)	Aqueous sodium hydroxide was added	to the first portion of the solution.
		observations	[1]
	(ii)		o the second portion of the solution. The mixture sted with a piece of filter paper soaked in aqueous plution.
		observations	
			[2]
(c)	Λfl	ame test was carried out on solid <b>E</b> .	
(0)			[4]
	ODS	Servations	[1]
tes	ts o	n solid F	
		tests	observations
The	solic	d was heated. The gas given off was	pungent gas evolved
		th damp, red litmus paper.	red litmus paper turned blue
			Tod mindo papor tarriod sido
		sodium hydroxide was added to	pungent gas evolved
		nd the mixture heated. The gas given ested.	Universal Indicator paper showed pH 10
(d)	Ide	ntify the gas given off in the tests on soli	id <b>F</b> . [1]
(e)	Ide	ntify <b>one</b> of the ions in solid <b>F</b> .	
			[1]
			[Total: 7]

Potassium suitate is the sait produced when suituric acid is neutralised by potassium hydroxide solution.
The correct amount of potassium hydroxide solution must be added to neutralise all of the sulfuric acid.
Plan an experiment to obtain pure crystals of potassium sulfate from sulfuric acid and potassium hydroxide solution.
You are provided with common laboratory apparatus.
[6]
[Total: 6]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2016 0620/62/M/J/16