

Cambridge IGCSE[™]

| CANDIDATE NAME | | | | | |
|-------------------|--|--|---------------------|--|--|
| CENTRE NUMBER | | | CANDIDATE NUMBER | | |

* 179218569

CHEMISTRY 0620/42

Paper 4 Theory (Extended)

February/March 2020

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS


- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

1 Petroleum is a useful natural resource.

The diagram shows how petroleum can be separated into useful substances.



| (a) | What is the name of the separation process shown in the diagram? | |
|-----|--|----|
| | | [2 |
| (b) | Name the fraction leaving at: | |
| | A | |
| | В | |
| | | [2 |
| (c) | Refinery gas is a mixture of hydrocarbons. | |

One refinery gas is butane, C₄H₁₀.

(i) Suggest the names of **two** other refinery gases.

..... and [2]

(ii) Write the chemical equation for the complete combustion of butane.


.....[2]

(iii) Name the toxic gas produced by the incomplete combustion of butane.

.....[1]

| (d) | Gas | soline and kerosene are both fuels. They have different properties. |
|-----|------|---|
| | (i) | Describe the differences in the properties given. |
| | | viscosity of the fuel |
| | | |
| | | flammability of the fuel |
| | | |
| | | |
| | (ii) | What difference in the molecules of gasoline and kerosene causes these differences in properties? |
| | | [1] |
| | | |
| (e) | Hyc | lrogen fuel cells can be used to power vehicles. |
| | Wri | te the word equation for the overall reaction that takes place in a hydrogen fuel cell. |
| | | [1] |
| | | [Total: 13] |

Aluminium is extracted from its ore. The ore is converted into pure aluminium oxide, which then undergoes electrolysis as shown.




| (a) | (i) | Name an ore of aluminium. | [4] |
|-----|-------|--|---------|
| | (ii) | What is meant by the term <i>electrolysis</i> ? | |
| | | | |
| (b) | | minium oxide has a melting point of about 2000 °C, but the electrolysis process operates out 900 °C. | s at |
| | (i) | Name the compound added to aluminium oxide to reduce the operating temperature. | [1] |
| | (ii) | Suggest one benefit to the environment of reducing the operating temperature. | |
| | | | |
| | (iii) | Write the ionic half-equation for the reaction taking place at: the negative electrode (cathode) | |
| | | the positive electrode (anode) | [4] |
| | (iv) | Explain why the anodes need frequent replacement. | |
| | | | |

| (c) Alu | minium oxide reacts with acids and with alkalis. |
|---------|---|
| (i) | What term is used to describe an oxide that reacts with acids and with alkalis? |
| (ii) | Aluminium oxide reacts with dilute sulfuric acid to form a salt. |
| () | State the name and write the formula of the salt formed. |
| | name |
| | formula[2] |
| (iii) | Aluminium oxide reacts with dilute sodium hydroxide to form a salt and one other product. |
| | Name the other product. |
| | [1] |
| (iv) | Aluminium hydroxide, $Al(OH)_3$, decomposes when heated to form aluminium oxide and water. |
| | Write the chemical equation for this reaction. |
| | [2] |
| (v) | Suggest the names of two other aluminium compounds that decompose when heated to form aluminium oxide. |
| | |
| | [2] |
| | [Total: 19] |

| The | The Periodic Table is a method of classifying elements. | | | |
|-----|---|------------------|------------------------|--|
| (a) | a) Identify the element which is in Group VI and Period 4. | | | |
| | | | | [1] |
| (b) | Calcium is in Group II an | d chlorine is in | n Group VII of the P | Periodic Table. |
| | Explain, in terms of num and chlorine atoms form | | | lectron transfer, how calcium atoms as formed. |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | [5] |
| (c) | (c) Group V chlorides are covalent molecules. The boiling points of some Group V chlorides are shown. | | | |
| | | chloride | boiling point/°C | |
| | | NCl ₃ | 71 | |
| | | PCl_3 | | |
| | | $AsCl_3$ | 130 | |
| | | $SbCl_3$ | 283 | |
| | (i) Suggest the approxi | mate boiling p | point of PCl_3 . | |
| | | | | [1] |
| | (ii) Explain the trend in | boiling points | in terms of attractive | e forces between particles. |
| | | | | [2] |
| | | | | |

(iii) Complete the dot-and-cross diagram to show the electron arrangement in a molecule of PCl_3 .

Show outer electrons only.



[3]

(d) PCl_3 reacts with chlorine, Cl_2 , to form PCl_5 . This reaction is exothermic and reaches an equilibrium.

$$PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g)$$

| (i) | Describe two features of an equilibrium. |
|-------|---|
| | |
| | [2 |
| (ii) | State the effect, if any, on the position of this equilibrium when the following changes are made. Explain your answers. |
| | temperature is increased |
| | |
| | pressure is increased |
| | [4 |
| (iii) | Explain, in terms of particles, what happens to the rate of the forward reaction when the reaction mixture is heated. |
| | |
| | |
| | |
| | |
| | |

| (e) PCl_5 reacts with lithium fluoride, LiF, to form LiPf | (e) | PCl ₅ reacts | with lithium | fluoride, | LiF, to | form | LiPF |
|---|-----|-------------------------|--------------|-----------|---------|------|------|
|---|-----|-------------------------|--------------|-----------|---------|------|------|

$$PCl_5 + 6LiF \rightarrow LiPF_6 + 5LiCl$$

Calculate the mass of LiF needed to form 3.04 g of ${\rm LiPF_6}$ using the following steps.

| • | Calculate the number of moles of LiPF ₆ formed. |
|---|--|
| | [M _r : LiPF ₆ , 152] |

| • | Deduce the number of moles of LiF needed. | number of moles = |
|---|---|-------------------|
| • | Calculate the mass of LiF needed. | number of moles = |

| mass = | | g |
|--------|----|----|
| | [3 | 31 |

- (f) Lithium fluoride has ionic bonding.
 - (i) What is an ionic bond?

| |
|------|
| [2] |

(ii) Give two physical properties of ionic compounds.



[Total: 28]

Iron is a typical transition element. 4

| ı | | _ | | |
|---|---|-----|---|--|
| ı | п | () | п | |

- acts as a catalyst forms coloured compounds
- has more than one oxidation state.

| (a) | | ne one major industrial process that uses iron as a catalyst and name the product made process. | in e |
|-----|-------|---|------|
| | prod | cess | |
| | prod | duct made | |
| | | | [2] |
| (b) | | en aqueous sodium hydroxide is added to aqueous $iron(II)$ sulfate, a precipitate forms. What colour is this precipitate? | |
| | (-) | | [4] |
| | (ii) | Write the ionic equation for this reaction. Include state symbols. | ניו |
| | | | [3] |
| (c) | | (II) sulfate can be converted to iron(III) sulfate by potassium manganate(VII) at rooperature. | om |
| | (i) | What is the role of potassium manganate(VII) in this reaction? | |
| | | | [1] |
| | (ii) | What condition must be used for this reaction to occur? | [-] |
| | | | [1] |
| (| (iii) | In terms of electron transfer, what happens to the $\text{iron}(II)$ ions in this reaction? | |
| | | | [1] |
| (| (iv) | State the colour change seen during this reaction. | |
| | | from purple to | [1] |
| | | | |
| (d) | Dec | luce the charge on the iron ion in each of these compounds. | |
| | FeF | 3 ····· | |
| | Fe(l | NO ₃) ₃ | |
| | | | [2] |

[Total: 12]

| 5 There are two | types of | polymers |
|-----------------|----------|----------|
|-----------------|----------|----------|

- (a) Addition polymers are made from many identical small units.
 - (i) What is the term used to describe these small units?

(ii) A section of an addition polymer is shown.

Draw the structure of the small unit used to make this addition polymer.


Show all of the atoms and all of the bonds.

(b) Polyamides are condensation polymers.
What does the term *condensation* mean when used to describe this type of polymer?


[2]

(c) A polyamide can be made from two different molecules.

A simplified structure of octanedioic acid is shown.



A simplified structure of 1,6-diaminohexane is shown.



| (i) | Complete the diagram to show a section of polyamide manufactured from octanedioic acid |
|-----|--|
| | and 1,6-diaminohexane. Include all of the atoms and all of the bonds in the linkages. |

| | | | | [3] |
|------|---------------------|-------------------|------|------------|
| (ii) | State the name of a | synthetic polyami | ide. | |
| | | | | [1] |
| | | | | [Total: 8] |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

| | ₹ | E 5 | helium 4 | 10 | Ne | neon 20 | 18 | Ā | argon 40 | 36 | 궃 | krypton 84 | 55 | Xe | xenon 131 | 98 | R | radon | | | |
|-------|---|-----|---------------|---------------|--------------|------------------------------|----|----|------------------|----|---------|-----------------|----|----------|------------------|-------|-------------|-----------------|--------|-----------|--------------------|
| | = | | | 6 | ட | fluorine 19 | 17 | Cl | chlorine 35.5 | 35 | ğ | bromine 80 | 53 | Н | iodine 127 | 85 | Αţ | astatine - | | | |
| | | | | 8 | 0 | oxygen 16 | 16 | S | sulfur 32 | 34 | Se | selenium 79 | 52 | <u>a</u> | tellurium 128 | 84 | Ъ | polonium — | 116 | ^ | livermorium - |
| | > | | | 7 | z | nitrogen 14 | 15 | ۵ | phosphorus 31 | 33 | As | arsenic 75 | 51 | Sb | antimony 122 | 83 | <u>B</u> | bismuth 209 | | | |
| | ≥ | | | 9 | O | carbon 12 | 14 | S | silicon 28 | 32 | Ge | germanium 73 | 20 | Sn | tin 119 | 82 | Pb | lead 207 | 114 | Εl | flerovium |
| | = | | | 5 | В | boron 11 | 13 | Ρl | aluminium 27 | 31 | Ga | gallium 70 | 49 | In | indium 115 | 81 | 11 | thallium 204 | | | |
| | | | | | | | | | | 30 | Zu | zinc 65 | 48 | ပ် | cadmium 112 | 80 | Ρ̈́ | mercury 201 | 112 | ű | copernicium |
| | | | | | | | | | | 29 | Cn | copper 64 | 47 | Ag | silver 108 | 62 | Au | gold 197 | 111 | Rg | roentgenium - |
| Group | | | | | | | | | | 28 | Z | nickel 59 | 46 | Pd | palladium 106 | 78 | ₽ | platinum 195 | 110 | Ds | darmstadtium - |
| Gre | | | | - | | | | | | 27 | ဝိ | cobalt 59 | 45 | 몬 | rhodium 103 | 77 | 'n | iridium 192 | 109 | ¥ | meitnerium - |
| | | - エ | hydrogen 1 | | | | | | | 26 | Fe | iron 56 | 44 | Ru | ruthenium 101 | 9/ | Os | osmium 190 | 108 | Hs | hassium - |
| | | | | | | | _ | | | 25 | Mn | manganese 55 | 43 | ပ | technetium - | 75 | Re | rhenium 186 | 107 | Bh | bohrium – |
| | | | | | pol | ass | | | | 24 | ပ် | chromium 52 | 42 | Mo | molybdenum 96 | 74 | ≥ | tungsten 184 | 106 | Sg | seaborgium - |
| | | | Key | atomic number | atomic symbo | name relative atomic mass | | | | 23 | > | vanadium 51 | 41 | Q N | niobium 93 | 73 | Б | tantalum 181 | 105 | Op | dubnium - |
| | | | | | atc | - re | | | | 22 | j | titanium 48 | 40 | Zr | zirconium 91 | 72 | Ξ | hafnium 178 | 104 | 짪 | rutherfordium - |
| | | | | | | | | | | 21 | Sc | scandium 45 | 39 | > | yftrium 89 | 57-71 | lanthanoids | | 89–103 | actinoids | |
| | = | | | 4 | Be | beryllium 9 | 12 | Mg | magnesium 24 | 20 | Ca | calcium 40 | 38 | ഗ് | strontium 88 | 56 | Ва | barium 137 | 88 | Ra | radium |
| | _ | | | 8 | = | lithium 7 | 1 | Na | sodium 23 | 19 | \prec | potassium 39 | 37 | & | rubidium 85 | 55 | Cs | caesium 133 | 87 | ቷ | francium - |
| | | | | | | | | | | | | | | | | | | | | | |

| _ | | | _ | _ | | | |
|-----|----|--------------|-----|-----|-----------|--------------|-----|
| 7.1 | Γn | lutetium | 175 | 103 | ۲ | lawrencium | I |
| 70 | Υp | ytterbium | 173 | 102 | 8 | nobelium | I |
| 69 | Tm | thulium | 169 | 101 | Md | mendelevium | 1 |
| 89 | Щ | erbinm | 167 | 100 | Fm | fermium | ı |
| 29 | 웃 | holmium | 165 | 66 | Es | einsteinium | I |
| 99 | Dy | dysprosium | 163 | 86 | ర్ | californium | I |
| 65 | Д | terbium | 159 | 26 | BK | berkelium | ı |
| 64 | Gd | gadolinium | 157 | 96 | Cm | curium | ı |
| 63 | En | europium | 152 | 92 | Am | americium | ı |
| 62 | Sm | samarium | 150 | 94 | Pn | plutonium | ı |
| 61 | Pm | promethium | I | 93 | Δ | neptunium | ı |
| 09 | pN | neodymium | 144 | 92 | \supset | uranium | 238 |
| 69 | Ā | praseodymium | 141 | 91 | Ра | protactinium | 231 |
| 58 | Ce | cerium | 140 | 06 | Ч | thorium | 232 |
| 22 | Гa | lanthanum | 139 | 88 | Ac | actinium | ı |

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).