



# Cambridge IGCSE™

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 9 5 3 4 6 3 7 5 2 1 \*



## CHEMISTRY

0620/43

Paper 4 Theory (Extended)

May/June 2023

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

### INSTRUCTIONS







- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

### INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.

This document has **16** pages.

1 Some symbol equations and word equations, **A** to **J**, are shown.



Use the equations to answer the questions that follow.

Each equation may be used once, more than once, or not at all.

Give the letter, **A** to **J**, for the equation which represents:

(a) photosynthesis ..... [1]

(b) an addition reaction ..... [1]

(c) a precipitation reaction ..... [1]

(d) incomplete combustion ..... [1]

(e) a displacement reaction ..... [1]

(f) a substitution reaction. .... [1]

[Total: 6]

**Question 2 starts on the next page.**

- 2 (a) The symbols of the elements in Period 3 of the Periodic Table are shown.

**Na    Mg    Al    Si    P    S    Cl    Ar**

Use the symbols of the elements in Period 3 to answer the questions that follow.  
Each symbol may be used once, more than once, or not at all.

Give the symbol of the element that:

- (i) is present in purified bauxite ..... [1]
- (ii) contains atoms with a full outer shell of electrons ..... [1]
- (iii) is used to kill microbes in water treatment ..... [1]
- (iv) forms an amphoteric oxide ..... [1]
- (v) forms an oxide which causes acid rain ..... [1]
- (vi) has an oxidation number of  $-1$  when it forms a compound with hydrogen.  
..... [1]
- (b) The relative atomic masses of elements can be calculated from the relative masses of isotopes and their percentage abundances.
- (i) Identify the isotope to which all relative masses are compared.  
..... [1]
- (ii) Table 2.1 shows the relative masses and the percentage abundances of the two isotopes in a sample of magnesium.

**Table 2.1**

relative mass of isotope	percentage abundance of isotope
24	85
26	15

Calculate the relative atomic mass of magnesium to **one** decimal place.

relative atomic mass = ..... [2]

(c) An ion contains 10 electrons, 13 protons and 14 neutrons.

(i) State the nucleon number of the ion.

..... [1]

(ii) Identify the element that forms this ion.

..... [1]

[Total: 11]

3 Magnesium forms ionic compounds.

- (a) Magnesium reacts with fluorine to form the ionic compound magnesium fluoride. The electronic configurations of an atom of magnesium and an atom of fluorine are shown in Fig. 3.1.




Fig. 3.1

- (i) Ions are formed by the transfer of electrons from magnesium atoms to fluorine atoms.

Complete the dot-and-cross diagrams in Fig. 3.2 to show the electronic configurations of **one** magnesium ion and **one** fluoride ion. Show the charges on the ions.




Fig. 3.2

[3]

- (ii) Deduce the formula of magnesium fluoride.

..... [1]

- (iii) When solid magnesium fluoride is dissolved in water it forms a solution that conducts electricity.

State one other change that can be made to solid magnesium fluoride to allow it to conduct electricity.

..... [1]

- (b) Silicon tetrachloride,  $\text{SiCl}_4$ , and silicon(IV) oxide,  $\text{SiO}_2$ , are covalent compounds.

Complete the dot-and-cross diagram in Fig. 3.3 to show the electronic configuration in a molecule of silicon tetrachloride. Show outer shell electrons only.




Fig. 3.3

[2]

- (c) The melting points of silicon tetrachloride and silicon(IV) oxide are shown in Table 3.1.

Table 3.1

	melting point/ $^{\circ}\text{C}$
silicon tetrachloride	-69
silicon(IV) oxide	1710

- (i) Silicon tetrachloride has a low melting point because it has weak forces of attraction between particles.

Name the type of particles that are held together by these weak forces of attraction.


..... [1]

- (ii) Explain, in terms of structure and bonding, why silicon(IV) oxide has a high melting point.

.....  
 .....  
 .....  
 ..... [2]

[Total: 10]

- 4 Hydrogen is produced by the reaction between zinc and dilute sulfuric acid,  $\text{H}_2\text{SO}_4$ .



- (a) A student carries out an experiment using excess zinc and dilute sulfuric acid.

The student measures the volume of hydrogen produced at regular time intervals using the apparatus shown in Fig. 4.1.

Lumps of zinc are used.




Fig. 4.1

The rate of reaction decreases as the reaction progresses. The rate eventually becomes zero.

- (i) Explain why the rate of reaction decreases as the reaction progresses.

.....  
 ..... [1]

- (ii) Explain why the rate of reaction eventually becomes zero.

.....  
 ..... [1]

- (b) The experiment is repeated using powdered zinc instead of lumps of zinc. All other conditions remain the same.

Explain, in terms of collision theory, why the rate of reaction increases if powdered zinc is used.

.....  
 .....  
 ..... [2]



(c) The equation for the reaction is shown.



25.0 cm<sup>3</sup> of 2.00 mol/dm<sup>3</sup> H<sub>2</sub>SO<sub>4</sub>(aq) is added to excess zinc.

Calculate the volume of H<sub>2</sub> formed at room temperature and pressure (r.t.p.). The volume of one mole of any gas is 24 dm<sup>3</sup> at r.t.p.

Use the following steps.

- Calculate the number of moles of H<sub>2</sub>SO<sub>4</sub> used.

..... mol

- Deduce the number of moles of H<sub>2</sub> produced.

..... mol

- Calculate the volume of H<sub>2</sub> formed at r.t.p.

..... dm<sup>3</sup>  
[3]

(d) Hydrogen can also be produced by the reaction of zinc with dilute hydrochloric acid.

- (i) Write a symbol equation for this reaction.

..... [2]

- (ii) State the test for hydrogen gas.

test .....

positive result .....

[1]

[Total: 10]

5 This question is about electricity and chemical reactions.

(a) Aqueous copper(II) sulfate is an electrolyte.

The electrolysis of aqueous copper(II) sulfate using inert electrodes forms:

- copper at the cathode
- oxygen at the anode.

(i) State what is meant by the term electrolyte.

.....  
 ..... [2]

(ii) State the term given to the Roman numeral, (II), in the name copper(II) sulfate.

..... [1]

(iii) State what happens to the colour of the aqueous copper(II) sulfate as this electrolysis progresses.

..... [1]

(iv) Write an ionic half-equation for the formation of copper at the cathode.

..... [2]

(v) Give the formula of the ion that forms oxygen at the anode.

..... [1]

(b) The electrolysis of aqueous copper(II) sulfate is repeated using **copper** electrodes.

State what happens to the anode.

..... [1]

(c) Spoons can be electroplated with silver.

(i) Name the substances used as:

the anode (positive electrode) .....

the cathode (negative electrode) .....

the electrolyte. ....

[3]

(ii) State **two** reasons why spoons are electroplated.

1 .....

2 .....

[2]

(d) Hydrogen–oxygen fuel cells can be used to produce electricity to power cars. Petrol produces carbon dioxide and carbon monoxide when it powers cars.

(i) State **one** adverse effect of carbon dioxide and carbon monoxide.

carbon dioxide .....

carbon monoxide ..... [2]


(ii) State **one** disadvantage, other than cost, of using hydrogen–oxygen fuel cells to power cars compared to using petrol.

..... [1]

[Total: 16]

6 This question is about nitrogen and compounds of nitrogen.

- (a) Ammonia is manufactured by the reaction between nitrogen and hydrogen in the Haber process.  
The equation is shown.



- (i) State the source of nitrogen for the Haber process.

..... [1]

- (ii) State the source of hydrogen for the Haber process.

..... [1]

- (iii) State the typical conditions used in the Haber process.

temperature ..... °C

pressure ..... atm

[2]

- (iv) Name the catalyst used in the Haber process.

..... [1]


- (v) State what is meant by the term catalyst.

.....

..... [2]

- (b) Ammonia is converted into nitric acid.

- (i) The first stage is the conversion of ammonia into nitrogen monoxide, NO.  
The equation is shown.



The reaction is carried out at a temperature of 900 °C and a pressure of 7 atm.  
The forward reaction is exothermic.

Using explanations that do **not** involve cost:

- explain why a temperature less than 900 °C is **not** used

.....


- explain why a pressure greater than 7 atm is **not** used.

.....

[2]

- (ii) In the second stage, nitrogen monoxide reacts with water and oxygen to produce nitric acid.

Balance the symbol equation for the reaction.



- (c) A student makes aqueous copper(II) nitrate by adding an excess of solid copper(II) carbonate to dilute nitric acid.

- (i) Write the symbol equation for this reaction.

..... [2]

- (ii) State **two** observations that indicate the copper(II) carbonate is in excess.

1 .....

2 .....

[2]


- (iii) Name **one** compound, other than copper(II) carbonate, that can be added to dilute nitric acid to produce aqueous copper(II) nitrate.

..... [1]

[Total: 15]

7 This question is about organic compounds.

(a) Propane and chlorine react at room temperature. An equation for the reaction is shown.



(i) State the condition required for this reaction.

..... [1]

(ii) Draw the displayed formulae of **two** structural isomers with the formula  $\text{C}_3\text{H}_7\text{Cl}$ .

[2]

(b) Alkenes are a homologous series of hydrocarbons.

(i) State **two** characteristics that all members of the same homologous series have in common.

1 .....

2 .....

[2]

(ii) Addition polymers are made from alkenes.

Complete Fig. 7.1 to show one repeat unit of the addition polymer formed from but-2-ene.




Fig. 7.1

[2]

- (c) A repeat unit of a condensation polymer is shown in Fig. 7.2.  
The polymer is made from two monomers.




Fig. 7.2

- (i) Draw the structures of the monomers used to produce the polymer in Fig. 7.2.

[2]

- (ii) Name the **type** of condensation polymer in Fig. 7.2.

..... [1]

- (iii) Name the **two** homologous series to which the monomers in (i) belong.

1 .....

2 .....

[2]

[Total: 12]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

## The Periodic Table of Elements

		Group															
I	II	III	IV	V	VI	VII	VIII										
1	2	3	4	5	6	7	8	9	10	11	12						
1	2	3	4	5	6	7	8	9	10	11	12						
H hydrogen 1	He helium 2	B boron 11	C carbon 12	N nitrogen 14	O oxygen 16	F fluorine 19	Ne neon 20	Al aluminium 13	Si silicon 28	P phosphorus 31	S sulfur 32	Cl chlorine 35.5	Ar argon 40				
<b>Key</b>																	
atomic number																	
atomic symbol																	
name																	
relative atomic mass																	
3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18		
Li lithium 7	Be beryllium 9	B boron 11	C carbon 12	N nitrogen 14	O oxygen 16	F fluorine 19	Ne neon 20	Na sodium 23	Mg magnesium 24	Al aluminium 27	Si silicon 28	P phosphorus 31	S sulfur 32	Cl chlorine 35.5	Ar argon 40		
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K potassium 39	Ca calcium 40	Sc scandium 45	Ti titanium 48	V vanadium 51	Cr chromium 52	Mn manganese 55	Fe iron 56	Co cobalt 59	Ni nickel 59	Cu copper 64	Zn zinc 65	Ga gallium 70	Ge germanium 73	As arsenic 75	Se selenium 79	Br bromine 80	Kr krypton 84
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb rubidium 85	Sr strontium 88	Y yttrium 89	Zr zirconium 91	Nb niobium 93	Mo molybdenum 96	Tc technetium —	Ru ruthenium 101	Rh rhodium 103	Pd palladium 106	Ag silver 108	Cd cadmium 112	In indium 115	Sn tin 119	Sb antimony 122	Te tellurium 128	I iodine 127	Xe xenon 131
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs caesium 133	Ba barium 137	lanthanoids	Hf hafnium 178	Ta tantalum 181	W tungsten 184	Re rhenium 186	Os osmium 190	Ir iridium 192	Pt platinum 195	Au gold 197	Hg mercury 201	Tl thallium 204	Pb lead 207	Bi bismuth 209	Po polonium —	At astatine —	Rn radon —
87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr francium —	Ra radium —	actinoids	Rf rutherfordium —	Db dubnium —	Sg seaborgium —	Bh bohrium —	Hs hassium —	Mt meitnerium —	Ds darmstadtium —	Rg roentgenium —	Cn copernicium —	Nh nihonium —	Fl flerovium —	Mc moscovium —	Lv livermorium —	Ts tennessine —	Og oganeson —

lanthanoids

actinoids

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La lanthanum 139	Ce cerium 140	Pr praseodymium 141	Nd neodymium 144	Pm promethium —	Sm samarium 150	Eu europium 152	Gd gadolinium 157	Tb terbium 159	Dy dysprosium 163	Ho holmium 165	Er erbium 167	Tm thulium 169	Yb ytterbium 173	Lu lutetium 175
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac actinium —	Th thorium 232	Pa protactinium 231	U uranium 238	Np neptunium —	Pu plutonium —	Am americium —	Cm curium —	Bk berkelium —	Cf californium —	Es einsteinium —	Fm fermium —	Md mendelevium —	No nobelium —	Lr lawrencium —

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).