## MARK SCHEME for the May/June 2014 series

## 0607 CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/22 Paper 2 (Extended), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



	Page 2	Mark Scheme	Syllabus	Paper		
		IGCSE – May/June 2014			0607	22
1	(a) ( (b) (	C plotted at $(5, -1)$		<b>B1</b> for eac	ch component	
	l,	7]				
2	(a)		1			
	(b)		2	<b>B1</b> for any shape with exactly 2 lines of symmetry		
	(c)		2	B1 for any order 4 bu each corn	y shape with rotatio at not all square or a er shaded	n symmetry of Ill 4 squares in
3	I 1	Labelled pie chart with angles 170, 100, 90	3	Tolerance <b>B1</b> for 1 a <b>B1</b> for 17	$2 \pm 2^{\circ}$ ingle drawn correctl 0, 100, 90 seen	y
4	(a) (	$0.29, 0.3, 33\%, \frac{1}{3} \sqrt{0.3}$	2	<b>B1</b> for 1 i	n wrong place	
	(b)	$\frac{\sqrt{5}}{2}, \frac{5}{\sqrt{5}}, 2\sqrt{5}, (\sqrt{5})^3$	2	<b>B1</b> for 1 i	n wrong place	
5	(a) 6	$5x^3 + 10x^3y$	2	<b>B1</b> for eac	ch term	
	(b) 2	$2a^2 - 7ab + 6b^2$	3	<b>B2</b> for 2 <i>a</i> <sup>2</sup> <b>B1</b> for abo	$a^2 - 4ab - 3ab + 6b^2$ ove with 1 wrong/c	mitted term
6	(a) (i) <sup>4</sup>	45	1	Accept in	factor form	
	(ii)	$\frac{4}{3}$	1	Accept $\frac{2^3}{3}$	2	
	(b) (i) 3	$3^{2} \times 5^{2}$	1			
	(ii) 2	$2^2 \times 3^3 \times 5^3 \times 7$	2	<b>B1</b> for 3 c	of 4 factors or <b>B1</b> fo	r 94 500

Page 3			Mark Scheme			Syllabus	Paper	
			IGCSE – May/June 2014			0607	22	
		-			-			
7	(a)	0.	3 0.25 0.3 0.4	2	M1 for di or B1 for	for dividing frequencies by number of trials, 1 for 3 correct		
	(b)	Μ	fore throws oe	1				
	(c)	64	10	1				
8	(a)	у	$=\frac{5\sqrt{x}}{2}$ oe	2	<b>M1</b> for <i>y</i> :	$=k\sqrt{x}$		
	(b)	$\frac{2}{2}$	$\frac{1}{5}$ oe	2	M1 for su	bstituting $y = 1$ in $t$	<i>heir</i> $y = k\sqrt{x}$	
9	(a)	4.	$9 \times 10^{-5}$	2	<b>M1</b> for 0. or 46 × 10	000046 + 0.000003 $)^{-6}$ or $0.3 \times 10^{-5}$		
	(b)	1.	$38  imes 10^{-10}$	2	<b>B1</b> for 13	$.8 \times 10^{-11}$		
10		(.:	$\frac{x-1}{(x+2)(2x+3)}$	3	M2 for $\frac{3}{-1}$ or M1 for	$\frac{(2x+3)-5(x+2)}{(x+2)(2x+3)}$ common denomina	ator $(x+2)(2x+3)$	