

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME			
 CENTRE NUMBER		CANDIDATE NUMBER	
PHYSICS Paper 6 Alterna	ative to Practical	0625/62 October/November 2013 1 hour	
	wer on the Question Paper. Iaterials are required.		

### READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.Write in dark blue or black pen.You may use a pencil for any diagrams or graphs.Do not use staples, paper clips, highlighters, glue or correction fluid.DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 12 printed pages and 4 blank pages.



1 The IGCSE class is investigating pendulums.

The apparatus is shown in Figs. 1.1 and 1.2.



[3]

For Examiner's Use (c) The student adjusts the length of the pendulum until its length l = 61.9 cm. He repeats For the procedure in (b) and calculates *T*. The results are shown in Table 1.1. Examiner's Use The student suggests that doubling the length l of the pendulum should double the period T. State whether the results support this suggestion and justify your answer by reference to the results. statement ..... justification ..... [2] (d) The student repeats the procedure in (b) four more times with different lengths of the pendulum. The student plots a graph of l against  $T^2$ . State two pieces of information from the graph that would indicate that *l* is directly proportional to  $T^2$ . 1. ..... 2. .... [2] (e) The student uses another pendulum. This pendulum has a mass that is double the mass of the first pendulum. Its length is 61.9 cm. The period T = 1.61 s. Suggest a conclusion about the effect of doubling the mass of the pendulum. .....[1] [Total: 9]


2 The IGCSE class is investigating the cooling of water.

A student cools some water by four different methods.

#### Experiment A (cooling with stirring)

(a) The student pours approximately  $200 \text{ cm}^3$  of hot water into a beaker.

She measures the temperature  $\theta_1$ . Fig. 2.1 shows the thermometer.



#### Fig. 2.1

Write down the temperature  $\theta_1$  shown on the thermometer in Fig. 2.1.

 $\theta_1 = \dots [1]$ 

For Examiner's Use

(b) The student stirs the water for one minute. She then records the temperature  $\theta_2$  of the water.

-10 0 10 20 30 40 50 60 70 80 90 100 110 

Fig. 2.2

(i) Write down the temperature  $\theta_2$  shown on the thermometer in Fig. 2.2.

*θ*<sub>2</sub> = .....

(ii) Calculate the temperature difference  $(\theta_1 - \theta_2)$ .

 $(\theta_1 - \theta_2) = \dots$ [1]

#### Experiment B (cooling with pouring)

(c) The student starts again with approximately  $200 \text{ cm}^3$  of hot water at the same initial temperature  $\theta_1$ .

She carefully pours the water from the beaker into another beaker. She pours the water back into the first beaker. She repeats this process four times.

She measures the temperature  $\theta_3$  of the water. Fig. 2.3 shows this temperature.

 $-10 \quad 0 \quad 10 \quad 20 \quad 30 \quad 40 \quad 50 \quad 60 \quad 70 \quad 80 \quad 90 \quad 100 \quad 110 \ ^{\circ}C$ 

Fig. 2.3

(i) Write down the temperature $\theta_3$ shown on the thermometer in Fig. 2.3.			
		$\theta_3 = \dots$	Examiner's Use
	(ii)	Calculate the temperature difference $(\theta_1 - \theta_3)$ .	
		$(\theta_1 - \theta_3) = \dots $ [1]	
Exp	oerin	nent C (cooling with a lid) and Experiment D (cooling without a lid)	
(d)		e student pours approximately 200 cm <sup>3</sup> of the hot water into each of two beakers. The al temperature of the water in each beaker is $\theta_1$ .	
	She	e places a lid on one of the beakers. She allows both beakers to cool for 5 minutes.	
	At t	he end of the cooling period, she calculates the temperature differences.	
		temperature difference of <b>C</b> (with a lid) = $\dots$ 11°C	
		temperature difference of <b>D</b> (without a lid) =	
		hk the experiments <b>A</b> , <b>B</b> , <b>C</b> and <b>D</b> in order, with the one that produced the greatest aperature drop first.	
		greatest temperature drop 1	
		2	
		3	
		smallest temperature drop 4[1]	
(e)		is laboratory investigation is to be repeated many times to check the results, suggest conditions that should be kept constant in order to provide reliable results.	
	1		
	2	[2]	
(f)	A st	tudent complains that the investigation is not a fair comparison.	
	Sug	ggest one way in which the investigation could be more fair.	
		[1]	
		[Total: 7]	

**3** The IGCSE class is investigating the resistance of a wire.

The circuit used is shown in Fig. 3.1.





Fig. 3.1

- (a) A student places the sliding contact **C** on the resistance wire **AB** at a distance *x* from **A**, where x = 0.200 m.
  - (i) He measures the current / in the wire. Fig. 3.2 shows the ammeter.





Record the value of *I*.

/ = .....[1]

For Examiner's

Use

(ii) The student measures the potential difference *V* across the wire between **A** and **C**. Fig. 3.3 shows the voltmeter.




Fig. 3.3

In the first row of Table 3.1 record the value of V.  $_{0625/62/O/N/13}$ 


(iii) Calculate the resistance *R* of the section **AC** of the wire using the equation  $R = \frac{V}{I}$ . Record *R* in the first row of the table.

<i>x</i> /m	V/V	$R/\Omega$
0.200		
0.350	0.80	2.67
0.500	1.00	3.33
0.650	1.25	4.17
0.800	1.60	5.33

## Table 3.1

(b) The student records the voltmeter readings using a range of *x* values. The readings are shown in Table 3.1.

Plot a graph of  $R/\Omega$  (y-axis) against x/m (x-axis).



[5]

[2]

For Examiner's Use (c) Using your graph, determine the length l of the resistance wire necessary to make a resistor of resistance 1.20  $\Omega$ . Show clearly on your graph how you obtained the necessary information.

For Examiner's Use

*l* = .....[1]


(d) Predict the resistance Z of 1.50 m of the resistance wire. Show your working.

*Z* = .....[1]

[Total: 10]


The IGCSE class is determining the focal length of a converging lens.			
Fig. 4.1 shows the apparatus.			
illuminated object u v flens Fig. 4.1			
<ul><li>(a) (i) On Fig. 4.1, measure and record the distance <i>u</i>, in mm, between the illuminated object and the lens.</li></ul>			
<i>u</i> = mm			
(ii) Measure and record the distance <i>v</i> , in mm, from the centre of the lens to the image on the screen.			
v =			
(iii) Calculate the value of <i>uv</i> .			
UV =			
(iv) Calculate the value of $(u + v)$ .			
$(u + v) = \dots$			
(v) Calculate a value $f_1$ for the focal length of the lens, using the equation $f_1 = \frac{uv}{(u+v)}$ .			
f <sub>1</sub> =[2]			
(b) A student does not move the position of the screen or the illuminated object. She moves the lens towards the screen until a smaller, sharply focused image of the object is seen on the screen.			
The new values of $u$ and $v$ are $u = \dots 42 \mathrm{mm}$			
v =			
(i) Calculate the value of <i>uv</i> .			
UV =			
(ii) Calculate the value of $(u + v)$ .			
$(u + v) = \dots$	I		


4



[1]

[Total: 9]





14

© UCLES 2013

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.