

Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME						
CENTRE NUMBER					CANDIDATE NUMBER	
BIOLOGY						9700/33
Advanced Practical Skills 1			Oc	tober/November 2014		
						2 hours
Candidates answer on the Question Paper.						
Additional Mate	erials:	As listed in th	ne Confic	dential Instructions.		

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
Total	



BLANK PAGE

Before you proceed, read carefully through the whole of Question 1 and Question 2.

Plan the use of your time to make sure that you finish all the work that you would like to do.

You will **gain marks** for recording your results according to the instructions.

If you have enough time, consider how you can improve the accuracy of your results, for example by obtaining and recording one or more additional measurements.

1 Plants translocate sap through phloem sieve tubes. Scientists discovered that aphids feed by pushing their stylet into a single phloem sieve tube as shown in Fig. 1.1.

By using stylets it is possible for scientists to collect the sap from a single phloem sieve tube.




Fig. 1.1

You are required to estimate, as accurately as possible, the concentration of sucrose in a sample of sap, **P**, by:

- carrying out a trial using the concentrations of sucrose solutions S1 and S2
- using the results of the trial to prepare further concentrations of sucrose solution
- obtaining more readings, so that you can estimate more accurately the concentration of sucrose in P.

A drop or small volume of **P** is placed into a known concentration of sucrose solution.

Fig 1.2 shows how the drop or small volume of **P** is released.





Fig. 1.2

Immediately the drop or small volume of **P** is released, the syringe is removed.

The drop of **P** may move up, move down or remain at the same level depending on the difference in the concentration between **P** and the sucrose solution.

(a) (i) Show clearly on the diagrams below, using the key, how you would expect to see the drop or small volume of **P** move if:



You are provided with:

labelled	contents	hazard	volume /cm³
S 1	0.10 mol dm ⁻³ sucrose solution	none	200
S 2	1.00 mol dm ⁻³ sucrose solution	none	200
Р	coloured sample of sap	none	20
W	distilled water	none	200

[1]

Read step 1 to step 8 before proceeding.

Proceed as follows:

You are required to estimate the concentration of **P** using **S1** and **S2**.

- 1. Place the 5 cm³ syringe on top of the large test-tube and use the marker pen to draw a line on the test-tube at the same height as the end of the syringe nozzle as shown in Fig. 1.2 (page 4).
- 2. Use the same syringe to suck up $4.0 \,\mathrm{cm}^3$ of **P**.
- 3. Put 35 cm³ of **S1** into the test-tube.
- 4. As shown in Fig.1.2, release one drop or a small volume of **P** and **remove** the syringe from the sucrose solution.

Note: **P** may come out without pushing the plunger, observe and record the direction of movement.

- 5. Immediately observe the movement of the drop or small volume of **P**.
- 6. Record your observations.
- 7. Wash out this test-tube.
- 8. Repeat steps 3 to 7 using **S2**.
 - (ii) Prepare the space below to record your observations.

(111)	Complete the following by using one of the words 'more' or 'less'.
	You may use each word once or more than once.
	P is concentrated than 0.1 mol dm ⁻³ (S1).
	P is concentrated than 1.0 mol dm ⁻³ (S2).
	Use your results to estimate the concentration of sucrose in P .
	estimate of concentration of sucrose in P[1]
(iv)	State which solution, S1 or S2, you will dilute to prepare further concentrations.
	Use your results to explain the reason for your decision.
	[2]

You are required to prepare further concentrations of sucrose solution by **simple** dilution, using the solution stated in **(a)(iv)**.

Decide on the further concentrations of sucrose solution you will use in your investigation.

You will need 40 cm³ of each sucrose solution.

- (v) Prepare the space below to show:
 - the concentration of each sucrose solution
 - the volumes of sucrose solution
 - the volumes of **W**.

(vi) Prepare the space below to record your observations.

9.	Prepare all the concentrations of sucrose solution you have listed in (a)(v) in the containers provided.
10.	Repeat steps 3 to 7 with each of the solutions.

(vii) Using these additional results from (vi) state a more accurate estimate of the concentration of sucrose in P than in (iii).

odify this procedure to investigate the concentration of solution.	Describe how you would mod sucrose in P , using Benedict's so	(viii)
[3]		
[Total: 17]		

You are required to use a sharp pencil for drawings and graphs.

Plants translocate sap through phloem sieve tubes. Scientists discovered that aphids feed by pushing their stylet into a single phloem sieve tube as shown in Fig. 2.1.
By using stylets it is possible for scientists to collect the sap from a single phloem sieve tube.




Fig. 2.1

Scientists have investigated the different rates of flow of sap in the stylets of aphids of different ages from hatching, \mathbf{H} , up to the adult, \mathbf{M} .

The results are shown in Table 2.1.

Table 2.1

different ages of aphid	rate of flow of sap /μl h ⁻¹			
	trial 1	trial 2	trial 3	mean
Н	0.452	0.448	0.451	0.450
J	0.726	0.722	0.726	0.725
K	0.977	1.232	0.973	
L	1.401	1.601	1.399	1.400
M	1.978	1.975	1.972	1.975


(a) (i) Circle the two anomalous results in Table 2.1.

[1]

(ii) Complete the table by calculating the missing mean value.

[1]

(iii) Plot a chart of the data in Table 2.1.



(iv)	Describe the relationship between the age of the aphid and the rate of flow of sap.	
		[1]
(v)	Suggest two reasons which might explain the relationship described in (iv).	
		[2]

[4]

(b) K1 is a slide of a stained transverse section through a plant stem. This plant species grows widely including Europe, Asia and North America.

You are not expected to be familiar with this specimen.

To help draw a plan diagram with the correct shape and proportions of the tissues, an eyepiece graticule can be used to measure the layers of tissues, without the need to calibrate the eyepiece graticule scale.

(i) Draw a large plan diagram of the part of the specimen on **K1** indicated by the shaded sector in Fig. 2.2.




Fig. 2.2

Use **one** ruled label line and label to show the pith.

[5]

(ii) Observe the tissue between the xylem and the hollow centre of the specimen.

Select one group of three cells with:


- one cell containing cell structures
- two cells which touch each other and also touch the cell which contains cell structures.

Make a large drawing of this group of three cells.

Use **one** ruled label line and label **D** to show **one** of the cell structures.

[5]

Fig. 2.3 is a photomicrograph of an aphid feeding on a plant.



magnification ×120

Fig. 2.3

(c) Calculate the actual length of the structure ${\bf R}$ to ${\bf T}$, using the magnification, giving your answer in μm .

You may lose marks if you do not show your working or if you do not use appropriate units.

actual lengthμm [4]

[Total: 23]

BLANK PAGE

BLANK PAGE

Copyright Acknowledgements:

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.