

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CHEMISTRY 9701/13

Paper 1 Multiple Choice May/June 2010

1 hour

Additional Materials: Multiple Choice Answer Sheet

Soft clean eraser

Soft pencil (type B or HB is recommended)

Data Booklet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Write your name, Centre number and candidate number on the Answer Sheet in the spaces provided unless this has been done for you.

There are **forty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A**, **B**, **C** and **D**.

Choose the one you consider correct and record your choice in soft pencil on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer. Any rough working should be done in this booklet.



Section A

For each question there are four possible answers, **A**, **B**, **C**, and **D**. Choose the **one** you consider to be correct.

- 1 Which gas closely approaches ideal behaviour at room temperature and pressure?
 - A ammonia
 - B carbon dioxide
 - C helium
 - **D** oxygen
- 2 Use of the Data Booklet is relevant to this question.

The elements radon (Rn), francium (Fr) and radium (Ra) have consecutive proton numbers in the Periodic Table.

What is the order of their first ionisation energies?

	least endothermic		most endothermic
Α	Fr	Ra	Rn
В	Fr	Rn	Ra
С	Ra	Fr	Rn
D	Rn	Ra	Fr

3 Use of the Data Booklet is relevant to this question.

What could be the proton number of an element that has three unpaired electrons in each of its atoms?

- **A** 5
- **B** 13
- **C** 15
- **D** 21

4 Given the following enthalpy changes,

$$I_2(g) + 3Cl_2(g) \rightarrow 2ICl_3(s)$$

$$\Delta H^{\circ} = -214 \,\mathrm{kJ} \,\mathrm{mol}^{-1}$$

$$I_2(s) \rightarrow I_2(g)$$

$$\Delta H^{\Theta} = +38 \,\mathrm{kJ} \,\mathrm{mol}^{-1}$$

What is the standard enthalpy change of formation of iodine trichloride, $ICl_3(s)$?

- A +176 kJ mol⁻¹
- **B** -88 kJ mol⁻¹
- $\mathbf{C} = 176 \,\mathrm{kJ} \,\mathrm{mol}^{-1}$
- **D** -214 kJ mol^{-1}

5 Some bond energy values are listed below.

bond	bond energy/kJ mol ⁻¹
C–H	410
C-C1	340
C <i>l</i> –C <i>l</i>	244
Br–Br	193

These bond energy values relate to the following four reactions.

P
$$Br_2 \rightarrow 2Br$$

Q
$$2Cl \rightarrow Cl_2$$

R
$$CH_3 + Cl \rightarrow CH_3Cl$$

S
$$CH_4 \rightarrow CH_3 + H$$

What is the order of enthalpy changes of these reactions from most negative to most positive?

A
$$P \rightarrow Q \rightarrow R \rightarrow S$$

B
$$Q \rightarrow R \rightarrow S \rightarrow P$$

$$\mathbf{C} \quad \mathsf{R} \to \mathsf{Q} \to \mathsf{P} \to \mathsf{S}$$

$$\mathbf{D} \quad \mathsf{S} \to \mathsf{P} \to \mathsf{Q} \to \mathsf{R}$$

Which mass of gas would occupy a volume of 3 dm³ at 25 °C and 1 atmosphere pressure? 6 [1 mol of gas occupies 24 dm³ at 25 °C and 1 atmosphere pressure.]

- **A** 3.2 g O₂ gas
- **B** 5.6 g N₂ gas
- C 8.0 g SO₂ gas
- **D** 11.0 g CO₂ gas

7 Ammonium nitrate, NH₄NO₃, can decompose explosively when heated.

$$NH_4NO_3 \rightarrow N_2O + 2H_2O$$

What are the changes in the oxidation numbers of the two nitrogen atoms in NH₄NO₃ when this reaction proceeds?

8 The Haber process for the manufacture of ammonia is represented by the following equation.

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

$$\Delta H = -92 \,\mathrm{kJ} \,\mathrm{mol}^{-1}$$

Which statement is correct about this reaction when the temperature is increased?

- A Both forward and backward rates increase.
- **B** The backward rate only increases.
- **C** The forward rate only increases.
- **D** There is no effect on the backward or forward rate.
- **9** Use of the Data Booklet is relevant to this question.

2.920 g of a Group II metal, \mathbf{X} , reacts with an excess of chlorine to form 5.287 g of a compound with formula $\mathbf{X}Cl_2$.

What is metal X?

- A barium
- **B** calcium
- **C** magnesium
- **D** strontium
- **10** The table gives the concentrations and pH values of the aqueous solutions of two compounds, X and Y. Either compound could be an acid or a base.

	Х	Y	
concentration	2 mol dm ⁻³	2 mol dm ⁻³	
рН	6	9	

Student P concluded that X is a strong acid.

Student Q concluded that the extent of dissociation is lower in X(aq) than in Y(aq).

Which of the students are correct?

- A both P and Q
- B neither P nor Q
- C Ponly
- **D** Q only

- 11 How does concentrated sulfuric acid behave when it reacts with sodium chloride?
 - A as an acid only
 - **B** as an acid and oxidising agent
 - C as an oxidising agent only
 - **D** as a reducing agent only
- **12** Swimming pool water can be kept free of harmful bacteria by adding aqueous sodium chlorate(I), NaOC*l*. This reacts with water to produce HOC*l* molecules which kill bacteria.

$$OCl^{-}(aq) + H_2O \rightleftharpoons OH^{-}(aq) + HOCl(aq)$$

In bright sunshine, the OCl⁻ ion is broken down by ultra-violet light.

$$OCl^{-}(aq) + uv light \rightarrow Cl^{-}(aq) + \frac{1}{2}O_{2}(g)$$

Which method would maintain the highest concentration of HOCl(aq)?

- A acidify the pool water
- B add a solution of chloride ions
- **C** add a solution of hydroxide ions
- **D** bubble air through the water
- 13 Na₂S₂O₃ reacts with dilute HCl to give a pale yellow precipitate. If 1 cm³ of 0.1 mol dm⁻³ HCl is added to 10 cm³ of 0.02 mol dm⁻³ Na₂S₂O₃ the precipitate forms slowly.

If the experiment is repeated with $1\,\mathrm{cm^3}$ of $0.1\,\mathrm{mol\,dm^{-3}}$ HC l and $10\,\mathrm{cm^3}$ of $0.05\,\mathrm{mol\,dm^{-3}}$ Na₂S₂O₃ the precipitate forms more quickly.

Why is this?

- **A** The activation energy of the reaction is lower when $0.05 \,\text{mol}\,\text{dm}^{-3}\,\text{Na}_2\text{S}_2\text{O}_3$ is used.
- **B** The reaction proceeds by a different pathway when $0.05\,\text{mol}\,\text{dm}^{-3}\,\text{Na}_2\text{S}_2\text{O}_3$ is used.
- **C** The collisions between reactant particles are more violent when 0.05 mol dm⁻³ Na₂S₂O₃ is used.
- **D** The reactant particles collide more frequently when 0.05 mol dm⁻³ Na₂S₂O₃ is used.

14 X is a salt of one of the halogens chlorine, bromine, iodine, or astatine (proton number 85).

The reaction scheme shows a series of reactions using a solution of X as the starting reagent.


$$X \xrightarrow{\mbox{HNO}_3(\mbox{aq})} \mbox{a precipitate} \xrightarrow{\mbox{an excess of} \\ \mbox{dilute NH}_3(\mbox{aq})} \mbox{a colourless} \\ \mbox{solution} \\ \mbox{an excess of} \\ \mbox{HNO}_3(\mbox{aq}) \\ \mbox{a precipitate}$$


What could X be?


- A sodium chloride
- B sodium bromide
- C potassium iodide
- **D** potassium astatide


15 The percentage of ammonia obtainable, if equilibrium were established during the Haber process, is plotted against the operating pressure for two temperatures, 400 °C and 500 °C.

Which diagram correctly represents the two graphs?









16 Two properties of non-metallic elements and their atoms are as follows.

property 1 has an oxide that can form a strong acid in water

property 2 has **no** paired 3p electrons

Which properties do phosphorus and sulfur have?

	phosphorus	sulfur	
Α	1 and 2	1 only	
В	1 only	1 and 2	
С	1 and 2	1 and 2	
D	2 only	only 1 only	

17 Consecutive elements X, Y, Z are in the third period of the Periodic Table. Element Y has the highest first ionisation energy and the lowest melting point.

What could be the identities of X, Y and Z?

- A aluminium, silicon, phosphorus
- B magnesium, aluminium, silicon
- C silicon, phosphorus, sulfur
- D sodium, magnesium, aluminium
- **18** Which property of Group II elements (beryllium to barium) decreases with increasing atomic number?
 - A reactivity with water
 - **B** second ionisation energy
 - C solubility of hydroxides
 - **D** stability of the carbonates
- 19 Which element of the third period requires the least number of moles of oxygen for the complete combustion of 1 mol of the element?
 - A aluminium
 - **B** magnesium
 - C phosphorus
 - **D** sodium

20 When gaseous chemicals are transported by road or by rail they are classified as follows.

flammable non-flammable poisonous

Which commonly transported gas is non-flammable?

- A butane
- **B** hydrogen
- C oxygen
- **D** propene

21 What will react differently with the two isomeric alcohols, (CH₃)₃CCH₂OH and (CH₃)₂CHCH₂CH₂OH?

- **A** acidified aqueous potassium manganate(VII)
- B concentrated sulfuric acid
- C phosphorus pentachloride
- **D** sodium

22 What is formed when propanone is refluxed with a solution of NaBH₄?

- A propanal
- B propan-1-ol
- C propan-2-ol
- **D** propane

23 When heated with chlorine, the hydrocarbon 2,2-dimethylbutane undergoes free radical substitution.

In a propagation step the free radical X* is formed.

How many different forms of X° are possible?

A 1

B 2

C 3

D 4

						9			
24	Wh	hich reagent will give similar results with both butanone and butanal?							
	Α	acidified aque	ous p	otassium dich	nromate	e(VI)			
	В	an alkaline sol	ution	containing co	omplexe	ed Cu ²⁺ ions (I	Fehling	g's solution)	
	С	an aqueous solution containing [Ag(NH ₃) ₂] ⁺ (Tollens' reagent)							
	D	2,4-dinitropher	nylhy	drazine reage	ent				
25		ich compound droxide?	is a	product of t	he hyd	rolysis of CH	l₃CO₂C	S₃H ₇ by boiling aqueous s	sodium
	A	CH ₃ OH	В	C ₃ H ₇ OH	С	$C_3H_7CO_2H$	D	$C_3H_7CO_2^-$ Na $^+$	
26	Wh	ich compound e	exhib	its both <i>cis-tra</i>	ans and	l optical isome	rism?		
	A	CH ₃ CH=CHCH	H ₂ CH	3					
	В	CH ₃ CHBrCH=	CH ₂						
	С	CH ₃ CBr=CBrC	CH ₃						
	D	CH ₃ CH ₂ CHBr(CH=C	CHBr					

27 In many countries plastic waste is collected separately and sorted. Some of this is incinerated to provide heat for power stations.

Why is pvc, polyvinylchloride, removed from any waste that is to be incinerated?

- **A** It destroys the ozone layer.
- B It does not burn easily.
- **C** It is easily biodegradable.
- **D** Its combustion products are harmful.
- 28 Polymerisation of 1,1-dichloroethene produces a dense, high melting point substance that does not allow gases to pass through. It is used as cling wrapping.

Which sequence appears in a short length of the polymer chain?

- A { CH₂CCl₂CH₂CCl₂CH₂CCl₂ }
- **B** {CHC/CHC/CHC/CHC/CHC/CHC/L
- C { CCl₂CCl₂CCl₂CCl₂CCl₂CCl₂CCl₂}
- **D** { CH₂CC*l*₂CHC*l*CHC*l*CH₂CC*l*₂ }

29 When an isomer Y of molecular formula C₄H₉Br undergoes hydrolysis in aqueous alkali to form an alcohol C₄H₉OH, the rate of reaction is found to be unaffected by changes in the concentration of OH⁻ ions present.

Which is the most likely molecular structure of Y?

- A CH₃CH₂CH₂CH₂Br
- B CH₃CH₂CHBrCH₃
- C (CH₃)₂CHCH₂Br
- D (CH₃)₃CBr
- **30** Which isomer of $C_4H_{10}O$ forms three alkenes on dehydration?
 - A butan-1-ol
 - **B** butan-2-ol
 - C 2-methylpropan-1-ol
 - **D** 2-methylpropan-2-ol

Section B

For each of the questions in this section, one or more of the three numbered statements 1 to 3 may be correct.

Decide whether each of the statements is or is not correct (you may find it helpful to put a tick against the statements that you consider to be correct).

The responses A to D should be selected on the basis of


A	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

- 31 Which reactions are redox reactions?
 - 1 CaBr₂ + 2H₂SO₄ \rightarrow CaSO₄ + Br₂ + SO₂ + 2H₂O
 - 2 CaBr₂ + 2H₃PO₄ \rightarrow Ca(H₂PO₄)₂ + 2HBr
 - 3 CaBr₂ + 2AgNO₃ \rightarrow Ca(NO₃)₂ + 2AgBr
- **32** Sodium hydrogensulfide, NaSH, is used to remove hair from animal hides.

Which statements about the SH⁻ ion are correct?

- 1 It contains 18 electrons.
- **2** Three lone pairs of electrons surround the sulfur atom.
- 3 Sulfur has an oxidation state of +2.
- 33 Which diagrams represent part of a giant molecular structure?



The responses A to D should be selected on the basis of

Α	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

34 In a car engine, non-metallic element X forms a pollutant oxide Y.

Further oxidation of Y to Z occurs in the atmosphere. In this further oxidation, 1 mol of Y reacts with $\frac{1}{2}$ mol of gaseous oxygen.

What can X be?

- 1 carbon
- 2 nitrogen
- 3 sulfur
- **35** When organic refuse decomposes in water carboxylic acids are formed. The water becomes acidic and aquatic life is destroyed.

Which additives are suitable to remove this acid pollution?

- 1 calcium carbonate
- 2 calcium hydroxide
- 3 potassium nitrate
- **36** Sulfur dioxide and sulfites are used in food preservation.


Why are they used for this purpose?

- 1 They are reducing agents so retard the oxidation of food.
- 2 They inhibit the growth of aerobic bacteria.
- 3 They react with $NO_2(g)$ converting it to NO(g).
- **37** Which structural formulae represent 2,2-dimethylpentane?
 - 1 $(CH_3)_2CHCH_2CH(CH_3)_2$
 - 2 (CH₃)₃CCH₂CH₂CH₃
 - 3 $CH_3CH_2CH_2C(CH_3)_3$

- 38 Which reactions are examples of nucleophilic substitution?
 - 1 $CH_3CH_2Br + OH^- \rightarrow CH_3CH_2OH + Br^-$
 - 2 $CH_3I + H_2O \xrightarrow{H^+} CH_3OH + HI$
 - 3 $CH_3CH_2CH_2Cl + NH_3 \rightarrow CH_3CH_2CH_2NH_2 + HCl$
- **39** Propanoic acid occurs naturally as a result of the bacterial fermentation of milk, and is partly responsible for the flavour of Swiss cheese.

Which starting materials could be used to synthesise propanoic acid?

- 1 CH₃CH₂CH₂OH
- 2 CH₃CH₂CN
- 3 CH₃CH₂CHO
- **40** The diagram shows some laboratory apparatus.



Which preparations could this apparatus be used for?

- 1 bromoethane, from ethanol, sodium bromide and concentrated sulfuric acid
- 2 ethanal, from ethanol, sodium dichromate(VI) and sulfuric acid
- 3 1,2-dibromoethane, from bromine and ethene

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.