

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CHEMISTRY 9701/11

Paper 1 Multiple Choice May/June 2013

1 hour

Additional Materials: Multiple Choice Answer Sheet

Soft clean eraser

Soft pencil (type B or HB is recommended)

Data Booklet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Write your name, Centre number and candidate number on the Answer Sheet in the spaces provided unless this has been done for you.

DO NOT WRITE IN ANY BARCODES.

There are **forty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A**, **B**, **C** and **D**.

Choose the one you consider correct and record your choice in soft pencil on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

Electronic calculators may be used.



Section A

For each question there are four possible answers, **A**, **B**, **C**, and **D**. Choose the **one** you consider to be correct.

1 Solutions containing chlorate(I) ions are used as household bleaches and disinfectants. These solutions decompose on heating as shown.

$$3ClO^- \rightarrow ClO_3^- + 2Cl^-$$

Which oxidation state is shown by chlorine in each of these three ions?

	C10 ⁻	C1O ₃ -	C <i>l</i> −
Α	+1	+3	-1
В	– 1	+3	+1
С	+1	+5	– 1
D	–1	+5	+1


2 A mixture of 10 cm³ of methane and 10 cm³ of ethane was sparked with an excess of oxygen. After cooling to room temperature, the residual gas was passed through aqueous potassium hydroxide.

All gas volumes were measured at the same temperature and pressure.

What volume of gas was absorbed by the alkali?

- **A** 15 cm³
- **B** 20 cm³
- \mathbf{C} 30 cm³
- \mathbf{D} 40 cm³

3 The diagram shows an electrolytic cell for the extraction of aluminium.



Which statement is correct?

A Aluminium ions are oxidised in this process.

B Aluminium is liberated at the anode by the reaction $Al^{3+} + 3e^{-} \rightarrow Al$.

C Cryolite is purified aluminium oxide.

D The graphite anode burns away.

4 Use of the Data Booklet is relevant to this question.

The elements radon (Rn), francium (Fr) and radium (Ra) have proton numbers 86, 87 and 88 respectively.

What is the order of their first ionisation energies?

	least endothermic		most endothermic
Α	Fr	Ra	Rn
В	Fr	Rn	Ra
С	Ra	Fr	Rn
D	Rn	Ra	Fr

- 5 In which species are the numbers of protons, neutrons and electrons all different?
 - $^{27}_{13}Al$
- **B** $^{35}_{17}\text{C}l^-$ **C** $^{32}_{16}\text{S}^{2-}$
- 6 An experiment is set up to measure the rate of hydrolysis of ethyl ethanoate.


$$CH_3CO_2C_2H_5 + H_2O \rightleftharpoons CH_3CO_2H + C_2H_5OH$$

The hydrolysis is found to be slow in neutral aqueous solution but it proceeds at a measurable rate when the solution is acidified with hydrochloric acid.

What is the function of the hydrochloric acid?

- to dissolve the ethyl ethanoate
- to ensure that the reaction reaches equilibrium В
- C to increase the reaction rate by catalytic action
- to suppress ionisation of the ethanoic acid formed

7 The reaction pathway for a reversible reaction is shown below.



Which statement is correct?

- The activation energy of the reverse reaction is +80 kJ mol⁻¹.
- The enthalpy change for the forward reaction is +30 kJ mol⁻¹. В
- C The enthalpy change for the forward reaction is +50 kJ mol⁻¹.
- The enthalpy change for the reverse reaction is +30 kJ mol⁻¹. D
- 8 Why does the rate of a gaseous reaction increase when the pressure is increased at a constant temperature?
 - More particles have energy that exceeds the activation energy.
 - В The particles have more space in which to move.
 - C The particles move faster.
 - D There are more frequent collisions between particles.
- 9 Which would behave the **least** like an ideal gas at room temperature?
 - carbon dioxide
 - В helium
 - C hydrogen
 - nitrogen
- **10** The general gas equation can be used to calculate the M_r value of a gas.

For a sample of a gas of mass mg, which expression will give the value of M_r ?

$$\mathbf{A} \quad M_{\rm r} = \frac{mpV}{RT}$$

$$\mathbf{C} \qquad M_{\rm r} = \frac{mR7}{pV}$$

A
$$M_r = \frac{mpV}{RT}$$
 B $M_r = \frac{pVRT}{m}$ **C** $M_r = \frac{mRT}{pV}$ **D** $M_r = \frac{pV}{mRT}$

11 A solution of Sn²⁺ ions will reduce an acidified solution of MnO₄⁻ ions to Mn²⁺ ions. The Sn²⁺ ions are oxidised to Sn⁴⁺ ions in this reaction.

How many moles of $\mathrm{Mn^{2+}}$ ions are formed when a solution containing 9.5 g of $\mathrm{SnC}l_2$ (M_r : 190) is added to an excess of acidified KMnO₄ solution?

- **A** 0.010
- **B** 0.020
- **C** 0.050
- **D** 0.125
- **12** Use of the Data Booklet is relevant to this question.

This question should be answered using bond enthalpy data. The equation for the complete combustion of methane is given below.

$$CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O$$

What is the enthalpy change of combustion of methane?

- **A** -1530 kJ mol⁻¹
- **B** -1184 kJ mol⁻¹
- $\mathbf{C} -770 \, \text{kJ} \, \text{mol}^{-1}$
- **D** $-688 \, \text{kJ} \, \text{mol}^{-1}$
- 13 In which row of the table are all statements comparing the compounds of magnesium and barium correct?

	solubility of	hydroxides	solubility of sulfates		
	solubility of magnesium hydroxide	solubility of barium hydroxide	solubility of magnesium sulfate	solubility of barium sulfate	
Α	higher	lower	higher	lower	
В	higher	lower	lower	higher	
С	lower	higher	higher	lower	
D	lower	higher	lower	higher	

- 14 What happens when iodine solution is added to a solution of sodium bromide?
 - **A** A reaction occurs without changes in oxidation state.
 - **B** Bromide ions are oxidised, iodine atoms are reduced.
 - **C** Bromide ions are reduced, iodine atoms are oxidised.
 - **D** No reaction occurs.

15 Element 85, astatine, is in Group VII. Concentrated sulfuric acid is added to sodium astatide. The mixture of products includes astatine, hydrogen astatide, hydrogen sulfide, and sodium sulfate.

Which product is formed by the oxidation of one of the constituents of sodium astatide?

- astatine Α
- hydrogen astatide В
- C hydrogen sulfide
- sodium sulfate D
- **16** Use of the Data Booklet is relevant to this question.

Magnesium nitrate, Mg(NO₃)₂, will decompose when heated to give a white solid and a mixture of gases. One of the gases released is an oxide of nitrogen, X.

7.4 g of anhydrous magnesium nitrate is heated until no further reaction takes place.

What mass of X is produced?

- **A** 1.5a
- **B** 2.3 g **C** 3.0 g
- **D** 4.6 g

17 Y is a salt of one of the halogens chlorine, bromine, iodine, or astatine (element 85).

The reaction scheme shows a series of reactions using a solution of Y as the starting reagent.

$$Y(aq) \xrightarrow{HNO_3(aq)} \text{ a precipitate } \xrightarrow{an \text{ excess of} \atop \text{dilute NH}_3(aq)} \text{ a colourless solution}$$

$$\downarrow \text{ an excess of} \atop \text{ an excess of} \atop \text{ HNO}_3(aq)$$

$$\text{ a precipitate}$$

What could Y be?

- A sodium chloride
- В sodium bromide
- C potassium iodide
- potassium astatide

18 Sulfur trioxide is manufactured from sulfur dioxide and oxygen, using the Contact process.

Which condition affects the value of the equilibrium constant, K_c ?

- A adjusting the temperature
- B increasing the pressure
- C removing SO₃ from the equilibrium mixture
- **D** using a catalyst
- 19 Which reagent, when mixed and heated with ammonium sulfate, liberates ammonia?
 - A aqueous bromine
 - **B** dilute hydrochloric acid
 - **C** limewater
 - **D** potassium dichromate(VI) in acidic solution
- **20** The following compounds are found in the seaweed *Asparagopsis taxiformis*.

Which compound could show **both** *cis-trans* isomerism and optical isomerism?

21 Lactic acid (2-hydroxypropanoic acid), CH₃CH(OH)CO₂H, is found in sour milk.

Which reaction could occur with lactic acid?

- A $CH_3CH(OH)CO_2H + CH_3OH \rightarrow CH_3CH(OCH_3)CO_2H + H_2O$
- **B** $CH_3CH(OH)CO_2H + HCO_2H \rightarrow CH_3CH(O_2CH)CO_2H + H_2O$
- **C** $CH_3CH(OH)CO_2H + NaHCO_3 \rightarrow CH_3CH(ONa)CO_2H + H_2O + CO_2$
- **D** $CH_3CH(OH)CO_2H + Cl_2 \rightarrow CH_3CH(Cl)CO_2H + HOCl$

22	Bro	mine reacts with	ethe	ene to form 1,2	-dibr	omoethane			
	Wh	hat is the correct description of the organic intermediate in this reaction?							
	Α	It has a negative	ch	arge.					
	В	It is a free radica	al.						
	С	It is a nucleophile	e.						
	D	It is an electroph	ile.						
23	Chl	oroethane can be	us	ed to make soc	dium	propanoate).		
				chloroethan	e →	$Q \rightarrow sod$	ium propa	anoate	
		e intermediate, C panoate.	Q, is	s hydrolysed v	with	boiling aqu	ieous so	dium hydroxide, to give	sodium
	Wh	ich reagent would	d pr	oduce the inter	medi	ate, Q, fron	n chloroet	hane?	
	Α	concentrated am	nmc	nia solution					
	В	dilute sulfuric aci	id						
	С	hydrogen cyanid	le						
	D	potassium cyani	de						
24	Aqı	ueous sodium hyd	drox	ide reacts with	1-bro	omopropan	e to give _l	propan-1-ol.	
	Hov	w should the first s	step	o in the mechar	nism	be describe	ed?		
	Α	by a curly arrow	froi	n a lone pair o	n the	OH ⁻ ion to	the C ^{δ+} at	tom of 1-bromopropane	
	В	by a curly arrow from the C ^{δ+} atom of 1-bromopropane to the OH ⁻ ion							
	С	by a curly arrow	froi	n the C–Br bor	nd to	the C atom			
	D	by the homolytic	fiss	sion of the C–B	r bor	nd			
25	fou							traviolet light. A compoundle contains of R contains of	
	Wh	ich two atoms of t	the	pentane chain	could	d be bonde	d to chlori	ne atoms in this isomer?	
	A	1 and 3	В	1 and 5	С	2 and 3	D	2 and 4	

- 26 Use of the Data Booklet is relevant to this question.
 - $2.30\,\mathrm{g}$ of ethanol were mixed with an excess of aqueous acidified potassium dichromate(VI). The reaction mixture was then boiled under reflux for one hour. The desired organic product was then collected by distillation. The yield of product was $60.0\,\%$.

What mass of product was collected?

- **A** 1.32 g
- **B** 1.38g
- **C** 1.80 g
- **D** 3.20 g
- 27 Burnt sugar has a characteristic smell caused partly by the following compound.

This compound contains two functional groups.

Which reagent will react with **only one** of the functional groups?

- **A** acidified potassium dichromate(VI)
- **B** 2,4-dinitrophenylhydrazine
- C hydrogen cyanide
- D sodium hydroxide
- **28** The structural formula of a compound **X** is shown below.

What is the name of compound \mathbf{X} and how does its boiling point compare with that of butanoic acid?

	name of X	boiling point of X
Α	methyl propanoate	higher
В	methyl propanoate	lower
С	propyl methanoate	higher
D	propyl methanoate	lower

29 Synthetic resins, plasticisers and many other chemicals can be made by polymerisation of a variety of monomers including prop-2-en-1-ol, CH₂=CHCH₂OH.

Which structure represents the repeat unit in poly(prop-2-en-1-ol)?

$$C - CH = C - I \\ CH_2OH$$

30 Some vegetable oils contain 'trans fats' that are associated with undesirable increases in the amount of cholesterol in the blood.

In the diagrams below, R₁ and R₂ are different hydrocarbon chains.

Which diagram correctly illustrates an optically active 'trans fat'?

Section B

For each of the questions in this section, one or more of the three numbered statements 1 to 3 may be correct.

Decide whether each of the statements is or is not correct (you may find it helpful to put a tick against the statements that you consider to be correct).

The responses A to D should be selected on the basis of

A	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

31 Use of the Data Booklet is relevant to this question.

Free-radicals play an important part in reactions involving the destruction of the ozone layer and the substitution of alkanes by chlorine.

Some free-radicals contain two unpaired electrons. Such species are called diradicals.

Which species are diradicals?

- 1 0
- 2 C1
- 3 CH₃

32 The Group II metals have higher melting points than the Group I metals.

Which factors could contribute towards the higher melting points?

- 1 There are smaller interatomic distances in the metallic lattices of the Group II metals.
- 2 More electrons are available from each Group II metal atom for bonding the atom into the metallic lattice.
- 3 Group II metals have a higher first ionisation energy than the corresponding Group I metal.
- 33 Valence shell electron pair repulsion theory should be used to answer this question.

Which species are trigonal planar?

- **1** BH₃
- 2 CH₃⁺
- 3 PH₃


The responses A to D should be selected on the basis of

Α	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

34 The first ionisation energies of twenty successive elements in the Periodic Table are represented in the graph.

The letters given are not the normal symbols for these elements.



Which statements about this graph are correct?

- 1 Elements *B*, *J* and *R* are in Group 0 of the Periodic Table.
- **2** Atoms of elements *D* and *L* contain two electrons in their outer shells.
- **3** Atoms of elements *G* and *O* contain a half-filled p subshell.

35 Solids W, X, Y and Z are compounds of two different Group II metals. Some of their applications are described below.

Compound **W** is used as a refractory lining material in kilns.

Compound **X** is used as a building material. It can also be heated in a kiln to form compound **Y**. When **Y** is hydrated, it forms compound **Z** which is used agriculturally to treat soils.

Which statements about these compounds are correct?

- More acid is neutralised by 1 g of W than by 1 g of X.
- 2 The metallic element in **W** reacts with water more quickly than the metallic element in **Y**.
- 3 Adding **Z** to a soil decreases the pH of the soil.
- 36 When a red-hot platinum wire is plunged into a test tube of hydrogen iodide, the gas is decomposed into its elements. If the experiment is repeated with hydrogen chloride, no change occurs.

Which factors contribute to this behaviour?

- 1 the strength of the hydrogen-halogen bond
- 2 the size of the halogen atom
- 3 the standard enthalpy of formation, ΔH_f^{\bullet} , of each of the products of decomposition
- **37** Which molecules would be present in the mixture produced by the photochemical chlorination of methane?
 - 1 hydrogen
 - 2 hydrogen chloride
 - 3 dichloromethane
- 38 In which reactions is the organic compound oxidised by the given reagent?
 - 1 CH₃CH₂CHO + Fehling's reagent
 - 2 CH₃CH₂CH₂CHO + Tollens' reagent
 - **3** CH₃CHO + 2,4-dinitrophenylhydrazine reagent

The responses A to D should be selected on the basis of

Α	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

39 A sun protection cream contains the following ester as its active ingredient.

Which substances are present in the products of its hydrolysis by aqueous sodium hydroxide?

1 CH₃CH₂CH₂CH₂CH(CH₂CH₃)CH₂OH

- **40** Which reagents, when used in an excess, can be used to make sodium lactate, $CH_3CH(OH)CO_2Na$, from lactic acid, $CH_3CH(OH)CO_2H$?
 - **1** Na
 - 2 NaHCO₃
 - 3 NaOH

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.