

# **Cambridge International Examinations**

Cambridge International Advanced Subsidiary and Advanced Level

CHEMISTRY 9701/13

Paper 1 Multiple Choice May/June 2017

1 hour

Additional Materials: Multiple Choice Answer Sheet

Soft clean eraser

Soft pencil (type B or HB is recommended)

Data Booklet

## **READ THESE INSTRUCTIONS FIRST**

Write in soft pencil.

Do not use staples, paper clips, glue or correction fluid.

Write your name, Centre number and candidate number on the Answer Sheet in the spaces provided unless this has been done for you.

DO **NOT** WRITE IN ANY BARCODES.

There are **forty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A**, **B**, **C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

## Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

Electronic calculators may be used.



International Examinations

CAMBRIDGE

# **Section A**

For each question there are four possible answers, A, B, C and D. Choose the one you consider to be correct.

Use of the Data Booklet may be appropriate for some questions.

1 The ion Y<sup>3-</sup> contains 18 electrons and has a mass number of 31.

How many protons and neutrons does Y<sup>3-</sup> contain?

	protons	neutrons
Α	15	16
В	15	18
С	18	13
D	21	10

A 0.216 g sample of an aluminium compound X reacts with an excess of water to produce a single hydrocarbon gas. This gas burns completely in O<sub>2</sub> to form H<sub>2</sub>O and CO<sub>2</sub> only. The volume of CO<sub>2</sub> at room temperature and pressure is 108 cm<sup>3</sup>.

What is the formula of X?

- $A Al_2C_3$
- **B**  $Al_3C_2$  **C**  $Al_3C_4$
- **D**  $Al_4C_3$
- 3 Which equation correctly describes the complete combustion of an alkene, C<sub>n</sub>H<sub>2n</sub>?

$$A \quad C_nH_{2n} \ + \ \tfrac{3}{2}\,nO_2 \ \rightarrow \ nCO_2 \ + \ 2nH_2O$$

$$\textbf{B} \quad C_nH_{2n} \ + \ \tfrac{3}{2}\,nO_2 \ \rightarrow \ nCO_2 \ + \ nH_2O$$

$$\mathbf{C} \quad \mathsf{C}_{\mathsf{n}}\mathsf{H}_{\mathsf{2n}} \,+\, \mathsf{2nO}_{\mathsf{2}} \,\rightarrow\, \mathsf{nCO}_{\mathsf{2}} \,+\, \mathsf{nH}_{\mathsf{2}}\mathsf{O}$$

$$\textbf{D} \quad C_nH_{2n} \ + \ 2nO_2 \ \rightarrow \ nCO_2 \ + \ 2nH_2O$$

4 Two conversions are shown.

$$NH_4^+ \rightarrow NH_3$$

$$C_2H_4 \rightarrow C_2H_6$$

Which similar feature do these two conversions have?

- A change in oxidation state of an element
- B decrease in bond angle
- **C** formation of a lone pair of electrons
- **D** loss of a  $\pi$  bond
- **5** All gases listed are at the same pressure.

Which gas will most closely approach ideal behaviour?

- A ammonia at 100 K
- **B** ammonia at 500 K
- C neon at 100 K
- **D** neon at 500 K
- **6** The following data are needed for this question.

$$\Delta H_{\rm f}^{\bullet}({\rm CO}(g)) = -111 \,\mathrm{kJ} \,\mathrm{mol}^{-1}$$

$$\Delta H_{f}^{\bullet}(CO_{2}(q)) = -394 \text{ kJ mol}^{-1}$$

$$\Delta H_f^{\bullet}(\text{Fe}_2\text{O}_3(\text{s})) = -822 \text{ kJ mol}^{-1}$$

Carbon monoxide reacts with iron(III) oxide.

$$3CO(g) + Fe_2O_3(s) \rightarrow 3CO_2(g) + 2Fe(s)$$

What is the enthalpy change when 55.8 g of iron are produced by this reaction?

- **A** –27.0 kJ
- **B** -13.5 kJ
- **C** +13.5 kJ
- **D** +27.0 kJ
- 7 Vanadium reacts with dilute sulfuric acid to form V<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> and hydrogen gas.

What happens to vanadium atoms in this reaction?

- **A** They lose three electrons and are oxidised.
- **B** They lose three electrons and are reduced.
- **C** They lose two electrons and are oxidised.
- **D** They lose two electrons and are reduced.

8 Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> reacts with HCl as shown.

$$Na_2S_2O_3(aq) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l) + SO_2(g) + S(s)$$

When calculating the oxidation number of sulfur in  $Na_2S_2O_3$ , the **average** oxidation number of the two sulfur atoms should be found.

What is the oxidation number of sulfur in each of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, SO<sub>2</sub>, and S?

	Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub>	SO <sub>2</sub>	S
Α	+2	+2	+1
В	+2	+4	0
С	+4	+4	0
D	+5	+4	0

**9** Ammonia is produced by the Haber process.

$$N_2 + 3H_2 \rightleftharpoons 2NH_3$$
  $\Delta H^{\circ} = -92 \text{ kJ mol}^{-1}$ 

A fault in the temperature control during the process resulted in the temperature changing to 600 °C for two hours.

What effect did this have on the ammonia production during this time?

- **A** Ammonia was formed faster. The equilibrium yield decreased.
- **B** Ammonia was formed faster. The equilibrium yield increased.
- **C** Ammonia was formed slower. The equilibrium yield decreased.
- **D** Ammonia was formed slower. The equilibrium yield increased.
- **10** For the reaction shown, an equilibrium is established at a temperature of 700 K.


The equilibrium constant,  $K_p$ , for the reaction is 9.80 kPa. The partial pressure of  $N_2O_4$  at equilibrium is 80.0 kPa.

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

What is the partial pressure of NO<sub>2</sub> at equilibrium?

**A** 8.16 kPa **B** 28.0 kPa **C** 66.6 kPa **D** 784 kPa

11 The diagram shows the Boltzmann distribution of the energy of gaseous molecules at a particular temperature.




Which statement is correct?


- **A** If the temperature of the gas is raised, the height of the maximum of the curve increases.
- **B** If the temperature of the gas is raised, the maximum of the curve moves to the right.
- **C** The length of the line labelled X shows the activation energy for the reaction.
- **D** The length of the line labelled Y shows the enthalpy change of the reaction.
- **12** Magnesium chloride, MgC $l_2$ , and silicon tetrachloride, SiC $l_4$ , are separately added to water.

What are the approximate pH values of the solutions formed?

	$MgC\mathit{l}_2$	SiCl <sub>4</sub>
Α	0–3	0–3
В	0–3	6–7
С	6–7	0–3
D	6–7	6–7

13 Which graph correctly shows the relative melting points of the elements Mg, A*l*, Si and P plotted against their relative first ionisation energies?





14 The table gives information about calcium carbonate and calcium hydroxide.

Which row is correct?

	calcium carbonate is more soluble than calcium hydroxide	calcium hydroxide can be manufactured using calcium carbonate as a starting material
Α	no	no
В	no	yes
С	yes	no
D	yes	yes

**15** Which row correctly describes the reactions of calcium and strontium with water?

	substance reduced	substance oxidised	more vigorous reaction
Α	calcium or strontium	water	calcium + water
В	calcium or strontium	water	strontium + water
С	water	calcium or strontium	calcium + water
D	water	calcium or strontium	strontium + water

**16** Chlorine gas is added to cold, aqueous sodium hydroxide.

In a separate experiment, chlorine gas is added to hot, aqueous sodium hydroxide.

Which oxidation states of chlorine are found in the reactants and products of the two reactions that take place?

- **A** 0, -1, +1 and +5
- **B** 0, –1 and +1 only
- **C** 0, –1 and +5 only
- **D** 0, +1 and +5 only

17 A test-tube of HBr(g) and a separate test-tube of HI(g) are heated to the same temperature.

Which combination of observations is possible?

		1
	test-tube of HBr(g)	test-tube of HI(g)
Α	a brown vapour appears	no change
В	a purple vapour appears	no change
С	no change	a brown vapour appears
D	no change	a purple vapour appears

**18** Ammonia exists as simple covalent molecules, NH<sub>3</sub>. Ammonia can react with suitable reagents to form products containing ammonium ions, NH<sub>4</sub><sup>+</sup>. Ammonia can also react with suitable reagents to form products containing amide ions, NH<sub>2</sub><sup>-</sup>.

Which of these nitrogen-containing species are present in an aqueous solution of ammonia?

- A ammonia molecules and amide ions only
- B ammonia molecules and ammonium ions only
- C ammonia molecules only
- **D** ammonium ions only

19			ld be pront				g of	nitroger	n mond	xide	react	with	an	excess	of	carbon
	Α	12 g of	carbon a	and	92 g of	nitrogen	diox	ide								
	В	24 g of	carbon a	and	92 g of	nitrogen	diox	ide								
	С	88 g of	carbon c	ixoib	de and	28 g of	nitrog	gen								
	D	88 g of	carbon c	ixoib	de and	56 g of	nitro	gen								
20	Stru	uctural i	somerisn	n an	d stere	oisomer	ism s	should b	e cons	idered	d whe	n ans	sweri	ng this	que	stion.
	Wh	ich form	nula ident	tifies	a <b>sing</b>	l <b>e</b> subs	tance	e?								
	Α	CH <sub>3</sub> CH	HC1CH₂C	НО												
	В	CH <sub>3</sub> CH	HCHCH <sub>3</sub>													
	С	CH <sub>2</sub> C1	CH <sub>2</sub> CHC	$:l_2$												
	D	C <sub>4</sub> H <sub>10</sub>														
21	Stru	uctural i	somerisn	n an	d stere	oisomer	ism s	should b	e cons	idered	d whe	n ans	weri	ng this	que	stion.
	If a	molecu	ıle contai	ns tv	wo non-	identica	ıl chir	al carb	on atom	ns, fou	ır opti	cal is	ome	rs exist.		
	Hov	w many	isomers	are	there w	vith										
		•	molecu	ular <sup>·</sup>	formula	C <sub>7</sub> H <sub>14</sub> C	and	I								
		•	a five-r	men	nbered	ring <b>and</b>	ł									
		•	a tertia	ary a	lcohol (	group?										
	Α	4		В	5		С	9		D	13					

22 The formulae of three compounds are shown.

C<sub>3</sub>H<sub>7</sub>CHO C<sub>2</sub>H<sub>5</sub>COCH<sub>3</sub> CH<sub>2</sub>CHCH<sub>2</sub>CH<sub>2</sub>OH

Only one of these compounds will decolourise bromine water. Only one of these compounds will produce a silver mirror with Tollens' reagent.

Which row shows the correct results?

	decolourises bromine water	forms a silver mirror with Tollens' reagent
Α	C₃H <sub>7</sub> CHO	C₂H₅COCH₃
В	C <sub>2</sub> H <sub>5</sub> COCH <sub>3</sub>	C₃H <sub>7</sub> CHO
С	CH <sub>2</sub> CHCH <sub>2</sub> CH <sub>2</sub> OH	C <sub>2</sub> H <sub>5</sub> COCH <sub>3</sub>
D	CH2CHCH2CH2OH	C <sub>3</sub> H <sub>7</sub> CHO

23 Carboxylic acids may be prepared by several different methods.

In which reaction would propanoic acid be formed?

- A adding ammonium propanoate to dilute sulfuric acid
- **B** heating ethyl propanoate with aqueous sodium hydroxide
- **C** heating propan-2-ol with acidified potassium manganate(VII) under reflux
- D heating propyl ethanoate with dilute sulfuric acid
- **24** The diagram represents the structure of a polymer.

By which method might this polymer be made?

- A polymerise ethene followed by hydration
- B polymerise ethene followed by oxidation with cold, acidified KMnO<sub>4</sub>
- **C** polymerise 1,2-dichloroethene followed by hydrolysis
- **D** polymerise 1,2-dichloroethene followed by oxidation with cold, acidified KMnO<sub>4</sub>

25 Bromine reacts with ethene in the dark.

Which description of the organic intermediate in this reaction is correct?

- A It has a negative charge.
- **B** It is a free radical.
- C It is a nucleophile.
- **D** It is an electrophile.
- 26 Which statement about butan-1-ol, butan-2-ol, and 2-methylbutan-2-ol is not correct?
  - A They all react with methanoic acid to form an ester.
  - **B** They all react with sodium.
  - **C** They can all be dehydrated to form an alkene.
  - **D** They can all be oxidised to a carbonyl compound.
- 27 Which compound will react with acidified potassium dichromate(VI) and with alkaline aqueous iodine?
  - A CH<sub>3</sub>COCH<sub>2</sub>CH<sub>3</sub>
  - B CH<sub>3</sub>CH(OH)CH<sub>2</sub>CH<sub>3</sub>
  - C CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CHO
  - D CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH

28 Chlorogenic acid is found in green coffee beans and is used in treatments for weight loss.

 $R = C_6H_5O_2$  and takes no part in the reaction with sodium carbonate.

chlorogenic acid

What is produced in good yield when chlorogenic acid is treated with an excess of sodium carbonate solution at room temperature?

Α

В

C

D

29 Butanoic acid can be produced from 1-bromopropane using reagents X and Y as shown.

What could be reagents X and Y?

	Х	Y
Α	KCN in ethanol	HC <i>l</i> (aq)
В	KCN in ethanol	NaOH(aq)
С	NH <sub>3</sub> in ethanol	HCℓ(aq)
D	NaOH(aq)	H <sup>+</sup> /Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup> (aq)

30 How many esters with the molecular formula  $C_5H_{10}O_2$  can be made by reacting a **primary** alcohol with a carboxylic acid?

**A** 4

**B** 5

**C** 6

**D** 8

#### **Section B**

For each of the questions in this section, one or more of the three numbered statements 1 to 3 may be correct.

Decide whether each of the statements is or is not correct (you may find it helpful to put a tick against the statements that you consider to be correct).

The responses A to D should be selected on the basis of

Α	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

Use of the Data Booklet may be appropriate for some questions.

- 31 Which statements about the atoms <sup>23</sup>Na and <sup>24</sup>Mg are correct?
  - 1 They have the same number of filled electron orbitals.
  - 2 They have the same number of neutrons.
  - **3** They are both reducing agents.
- **32** Which compounds contain covalent bonds?
  - 1 aluminium chloride
  - 2 ammonia
  - 3 calcium fluoride
- **33** Ethylene glycol, HOCH<sub>2</sub>CH<sub>2</sub>OH, is used as a de-icer. It allows ice to melt at temperatures below 0 °C.

Which statements are correct?

- 1 Ethylene glycol changes the extensive network of hydrogen bonds in ice.
- 2 Ethylene glycol molecules form hydrogen bonds with other ethylene glycol molecules.
- **3** Ethylene glycol molecules will dissolve in the water formed from the ice.

The responses A to D should be selected on the basis of

Α	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

**34** A student carried out two experiments using MnO<sub>2</sub> as a catalyst to decompose hydrogen peroxide. The equation for this reaction is shown.

$$2H_2O_2(aq) \rightarrow 2H_2O(I) + O_2(g)$$

The student's results are recorded in the table.

	mass MnO <sub>2</sub>	conditions	vol H <sub>2</sub> O <sub>2</sub>	final vol O <sub>2</sub>	time taken
experiment 1	0.25 g	room conditions	10.0 cm <sup>3</sup>	480 cm <sup>3</sup>	200 s
experiment 2	0.25 g	room conditions	10.0 cm <sup>3</sup>	480 cm <sup>3</sup>	500s

Which statements are correct?

- 1 The activation energy was the same for both experiments.
- 2 The concentration of the hydrogen peroxide solution used was 4.0 mol dm<sup>-3</sup>.
- 3 The MnO<sub>2</sub> used in experiment 1 was in larger pieces than the MnO<sub>2</sub> used in experiment 2.
- 35 The formula of a solid magnesium compound is Mg(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O. When the solid compound is heated in a hard-glass test-tube, one or more changes are observed.

Which statements are correct?

- 1 Strong heating produces a basic residue.
- 2 Strong heating produces a brown gas.
- **3** Gentle heating produces colourless droplets of liquid on the upper parts of the test-tube.
- **36** Sulfur dioxide is an atmospheric pollutant.

What might result from the release of sulfur dioxide gas into the atmosphere?

- 1 reduction of NO<sub>2</sub> to NO
- 2 depletion of aquatic life
- 3 corrosion of limestone statues

37 Which mechanisms are represented in this sequence of reactions?

$$CH_3CH_2CH_3 \rightarrow CH_3CH_2CH_2Cl \rightarrow CH_3CH_2CH_2OH \rightarrow CH_3CHCH_2 \rightarrow CH_3CHBrCH_2Br$$

- 1 electrophilic addition
- 2 nucleophilic substitution
- 3 free radical substitution
- 38 Which compounds, on heating with ethanolic NaOH, produce more than one product with molecular formula C₄H<sub>8</sub>?
  - 1 2-bromobutane
  - 2 2-bromo-2-methylpropane
  - 3 1-bromo-2-methylpropane
- **39** Substance M is refluxed with aqueous sodium hydroxide. One of the products of this reaction reacts with alkaline aqueous iodine to give a pale yellow precipitate.

Which compounds could be substance M?

- 1 CH<sub>3</sub>CO<sub>2</sub>CH<sub>3</sub>
- 2 CH<sub>3</sub>CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>
- 3 HCO<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>
- **40** Propanal will react with hydrogen cyanide to form 2-hydroxybutanenitrile. A suitable catalyst for this reaction is sodium cyanide.

$$\begin{array}{c} \text{NaCN} \\ \text{CH}_3\text{CH}_2\text{CHO} + \text{HCN} & \Longrightarrow & \text{CH}_3\text{CH}_2\text{CH(OH)CN} \end{array}$$

Which statements about this reaction of propanal with hydrogen cyanide are correct?

- 1 The CN<sup>-</sup> ion attacks the propanal molecule to form an intermediate ion.
- 2 The product of the reaction has a chiral carbon atom.
- **3** The CN<sup>-</sup> ion is a stronger electrophile than the HCN molecule.

# **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.