



Cambridge International AS & A Level

CANDIDATE NAME



CENTRE NUMBER

--	--	--	--	--

CANDIDATE NUMBER

--	--	--	--



CHEMISTRY

9701/53

Paper 5 Planning, Analysis and Evaluation

October/November 2024

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 30.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages. Any blank pages are indicated.





1 A student uses the following method to determine the percentage by mass of the painkiller aspirin, $C_9H_8O_4(s)$, in some tablets.

step 1 Grind five tablets into a powder.

step 2 Use a weighing boat to accurately weigh by difference approximately 0.4 g of powdered tablets into a pear-shaped flask containing anti-bumping granules.

step 3 Add 25cm^3 of aqueous 1mol dm^{-3} sodium hydroxide, NaOH(aq) , to the pear-shaped flask, forming mixture **A**.

step 4 Reflux mixture **A** for 20 minutes.

step 5 Allow mixture **A** to cool and then filter into a small beaker. Label the filtrate solution **B**.

step 6 Add 30cm^3 of alkaline aqueous iodine to solution **B** and leave to stand for 1 hour. A precipitate, **C**, $(C_6H_2I_2O)_2(s)$, will form.

step 7 Filter the resulting mixture under reduced pressure. Wash the residue, **C**, with a small volume of cold distilled water.

step 8 Allow solid **C** to dry.

step 9 Weigh solid **C** and record its mass.

Alkaline aqueous iodine is irritating to the skin and eyes.

(a) Identify an appropriate precaution, other than eye protection and a lab coat, that the student should take when using alkaline aqueous iodine.

.....
..... [1]

(b) Describe how the student should carry out **step 2**. Include a results table, with appropriate headings, for the student to fill in.

.....
.....
.....

[2]





- (c) Complete Fig. 1.1 to show how **step 4** is carried out in the laboratory.
Label your diagram fully.




Fig. 1.1

[2]

- (d) (i) The student uses a measuring cylinder to measure the volume of alkaline aqueous iodine in **step 6**. Suggest why this is a suitable piece of apparatus to use.

.....
 [1]

- (ii) Suggest why the student leaves the mixture to stand for 1 hour in **step 6**.

.....
 [1]

- (iii) Explain why the residue is washed in **step 7**.

.....
 [1]

- (iv) Explain why hot distilled water is **not** used in **step 7**.


.....
 [1]




DO NOT WRITE IN THIS MARGIN



- (e) The equation for the reaction between aspirin, $C_9H_8O_4(s)$, and $NaOH(aq)$, which takes place in **step 4**, is shown.



The equation for the reaction in which solid **C**, $(C_6H_2I_2O)_2(s)$, is formed in **step 6** is shown.




The student's results are shown in Table 1.1.

Table 1.1

mass of powdered tablets added to the pear-shaped flask in step 2	0.409 g
mass of dry $(C_6H_2I_2O)_2(s)$ recorded in step 9	0.764 g

- (i) Calculate the amount, in mol, of $(C_6H_2I_2O)_2(s)$ collected in **step 9**.



amount of $(C_6H_2I_2O)_2$ mol [1]

- (ii) Use your answer to (i) to calculate the mass, in g, of $C_9H_8O_4(s)$ in the powdered tablets added to the flask in **step 2**.

mass of $C_9H_8O_4(s)$ g [1]

- (iii) Use your answer to (ii) to calculate the percentage by mass of aspirin, $C_9H_8O_4(s)$, in the tablets.

If you were unable to obtain an answer to (ii) you may use 0.374 g for the mass of $C_9H_8O_4(s)$. This is **not** the correct value.

percentage by mass $C_9H_8O_4(s)$ in the tablets [1]





(f) Another student follows the same method but does not allow solid **C** to dry completely in **step 8**.

State and explain the effect that this has on the calculated percentage by mass of aspirin, $C_9H_8O_4(s)$, in the tablets.

.....
.....
..... [1]

[Total: 13]

DO NOT WRITE IN THIS MARGIN





2 Crystal violet, $C_{25}H_{30}N_3Cl(s)$, is a purple dye.

Some light is absorbed when it passes through $C_{25}H_{30}N_3Cl(aq)$.

Absorbance is the proportion of light absorbed at a particular wavelength. This is measured using a colorimeter.

A graph of absorbance against wavelength for $C_{25}H_{30}N_3Cl(aq)$ is shown in Fig. 2.1.




Fig. 2.1

A student investigates how to determine the concentration of aqueous crystal violet, $C_{25}H_{30}N_3Cl(aq)$, using colorimetry.

(a) Suggest the best wavelength of light to use in the colorimeter when measuring the concentration of $C_{25}H_{30}N_3Cl(aq)$.

wavelength =nm [1]

DO NOT WRITE IN THIS MARGIN





(b) Solution D is 500.0 cm³ of 2.50 × 10⁻² mol dm⁻³ C₂₅H₃₀N₃Cl(aq).

(i) Calculate the mass of C₂₅H₃₀N₃Cl(s) needed to prepare solution D.

Give your answer to **three significant figures**.

[M_r: C₂₅H₃₀N₃Cl(s), 407.5]

mass of C₂₅H₃₀N₃Cl(s) =g [1]

(ii) The student is given a small beaker containing the mass of C₂₅H₃₀N₃Cl(s) calculated in (i).

Describe how the student should prepare 500.0 cm³ of solution D.

Include the name and capacity of the key apparatus which should be used and describe how the student should ensure the volume is exactly 500.0 cm³.

.....

.....

.....

.....

.....

.....

.....

..... [3]

DO NOT WRITE IN THIS MARGIN





- (c) A small sample of solution **D** was diluted to form solution **E**, $2.50 \times 10^{-4} \text{ mol dm}^{-3}$ $\text{C}_{25}\text{H}_{30}\text{N}_3\text{Cl}(\text{aq})$.

The student prepares solutions 2 to 6 as shown in Table 2.1.

The total volume needed for each of solutions 2 to 6 is 20.00 cm^3 .

Each solution is placed into a colorimeter and the absorbance is measured.

- (i) Complete Table 2.1 to show the volumes of solution **E** and distilled water needed to prepare each of the solutions from 2 to 6. Give all volumes to **two** decimal places.

Table 2.1

solution	volume of $2.50 \times 10^{-4} \text{ mol dm}^{-3}$ $\text{C}_{25}\text{H}_{30}\text{N}_3\text{Cl}(\text{aq})$ (solution E) $/ \text{cm}^3$	volume of distilled water $/ \text{cm}^3$	$[\text{C}_{25}\text{H}_{30}\text{N}_3\text{Cl}(\text{aq})]$ $/ \text{mol dm}^{-3}$	absorbance
1	0.00	20.00	0.00	0.000
2			0.50×10^{-4}	0.191
3			1.00×10^{-4}	0.270
4			1.50×10^{-4}	0.545
5			2.00×10^{-4}	0.711
6			2.50×10^{-4}	0.860

[1]

- (ii) Identify the dependent variable.

..... [1]





(d) (i) Plot a graph of absorbance against $[C_{25}H_{30}N_3Cl(aq)]$ on the grid in Fig. 2.2.

Use a cross (x) to plot each data point.

Draw a straight line of best fit.




Fig. 2.2



DO NOT WRITE IN THIS MARGIN



(ii) Circle the point on the graph you consider to be most anomalous.

Suggest **one** reason why this anomaly may have occurred during this experimental procedure.

Assume no error was made in the measurement of absorbance.

.....
..... [2]

(iii) State the relationship between $[C_{25}H_{30}N_3Cl(aq)]$ and absorbance.

..... [1]

(iv) Suggest how the student could improve the reliability of the data obtained in the experiment in (c).

.....
..... [1]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN






Question 2 continues on the next page.



DO NOT WRITE IN THIS MARGIN



- (e) The student carries out a further experiment to examine the kinetics of the reaction between crystal violet, $C_{25}H_{30}N_3Cl(aq)$, and aqueous sodium hydroxide, $NaOH(aq)$.



The disappearance of the purple colour as the reaction proceeds can be monitored by measuring how the absorbance of light by the mixture changes using a colorimeter.

The student mixes 5 cm^3 of solution 6 with 5 cm^3 of $NaOH(aq)$, a large excess, and immediately starts the stopwatch.

The resulting mixture is then placed in a colorimeter. The absorbance of this mixture is measured every 100 seconds after starting the stop-watch.

Fig. 2.3 shows a graph of the student's results.




Fig. 2.3





(i) Suggest why it is **not** possible for the student to measure the absorbance of the mixture at $t = 0$ s.

.....
..... [1]

(ii) Use the graph in Fig. 2.3 to find the half-life, $t_{1/2}$, starting at 100 s.

State the coordinates of both points on the line of best fit used in your calculation.

coordinates 1 coordinates 2
half-life s [2]

(iii) Another student repeats the experiment at a different temperature and measures two half-life values. The values obtained are 420 s and 425 s.

Use these values to deduce the order of the reaction with respect to $C_{25}H_{30}N_3Cl(aq)$. Explain your answer.

order =
explanation [1]

[Total: 17]

DO NOT WRITE IN THIS MARGIN





BLANK PAGE

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$)





The Periodic Table of Elements

		Group																																																																								
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18																																																									
		<table border="1" style="margin: auto;"> <tr> <td>1</td> <td>H</td> <td>hydrogen</td> <td>1.0</td> </tr> </table>																1	H	hydrogen	1.0																																																					
1	H	hydrogen	1.0																																																																							
		<table border="1" style="margin: auto;"> <tr> <td colspan="2" style="text-align: center;">Key</td> </tr> <tr> <td style="text-align: center;">atomic number</td> <td style="text-align: center;">atomic symbol</td> </tr> <tr> <td style="text-align: center;">name</td> <td style="text-align: center;">relative atomic mass</td> </tr> </table>																Key		atomic number	atomic symbol	name	relative atomic mass																																																			
Key																																																																										
atomic number	atomic symbol																																																																									
name	relative atomic mass																																																																									
3	Li	Li	lithium	6.9	4	Be	beryllium	9.0	21	Sc	scandium	44.9	22	Ti	titanium	47.9	23	V	vanadium	50.9	24	Cr	chromium	52.0	25	Mn	manganese	54.9	26	Fe	iron	55.8	27	Co	cobalt	58.9	28	Ni	nickel	58.7	29	Cu	copper	63.5	30	Zn	zinc	65.4	31	Ga	gallium	69.7	32	Ge	germanium	72.6	33	As	arsenic	74.9	34	Se	selenium	79.0	35	Br	bromine	79.9	36	Kr	krypton	83.8		
11	Na	Na	sodium	23.0	12	Mg	magnesium	24.3	39	Y	yttrium	88.9	40	Zr	zirconium	91.2	41	Nb	niobium	92.9	42	Mo	molybdenum	95.9	43	Tc	technetium	—	44	Ru	ruthenium	101.1	45	Rh	rhodium	102.9	46	Pd	palladium	106.4	47	Ag	silver	107.9	48	Cd	cadmium	112.4	49	In	indium	114.8	50	Sn	tin	118.7	51	Sb	antimony	121.8	52	Te	tellurium	127.6	53	I	iodine	126.9	54	Xe	xenon	131.3		
19	K	K	potassium	39.1	20	Ca	calcium	40.1	57–71	lanthanoids					72	Hf	hafnium	178.5	73	Ta	tantalum	180.9	74	W	tungsten	183.8	75	Re	rhenium	186.2	76	Os	osmium	190.2	77	Ir	iridium	192.2	78	Pt	platinum	195.1	79	Au	gold	197.0	80	Hg	mercury	200.6	81	Tl	thallium	204.4	82	Pb	lead	207.2	83	Bi	bismuth	209.0	84	Po	polonium	—	85	At	astatine	—	86	Rn	radon	—
37	Rb	Rb	rubidium	85.5	38	Sr	strontium	87.6	89–103	actinoids					104	Rf	rutherfordium	—	105	Db	dubnium	—	106	Sg	seaborgium	—	107	Bh	bohrium	—	108	Hs	hassium	—	109	Mt	meitnerium	—	110	Ds	darmstadtium	—	111	Rg	roentgenium	—	112	Cn	copernicium	—	113	Nh	nihonium	—	114	Fl	flerovium	—	115	Mc	moscovium	—	116	Lv	livermorium	—	117	Ts	tennessine	—	118	Og	oganeson	—

lanthanoids

actinoids

57	La	La	lanthanum	138.9	58	Ce	cerium	140.1	59	Pr	praseodymium	140.9	60	Nd	neodymium	144.2	61	Pm	promethium	—	62	Sm	samarium	150.4	63	Eu	europium	152.0	64	Gd	gadolinium	157.3	65	Tb	terbium	158.9	66	Dy	dysprosium	162.5	67	Ho	holmium	164.9	68	Er	erbium	167.3	69	Tm	thulium	168.9	70	Yb	ytterbium	173.1	71	Lu	lutetium	175.0
89	Ac	Ac	actinium	—	90	Th	thorium	232.0	91	Pa	protactinium	231.0	92	U	uranium	238.0	93	Np	neptunium	—	94	Pu	plutonium	—	95	Am	americium	—	96	Cm	curium	—	97	Bk	berkelium	—	98	Cf	californium	—	99	Es	einsteinium	—	100	Fm	fermium	—	101	Md	mendeleevium	—	102	No	nobelium	—	103	Lr	lawrencium	—

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

