



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 1 3 6 0 5 2 4 7 3 1 *

MATHEMATICS

9709/13

Paper 1 Pure Mathematics 1

May/June 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.


INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

- 3 In each of parts (a), (b) and (c), the graph shown with solid lines has equation $y = f(x)$. The graph shown with broken lines is a transformation of $y = f(x)$.

(a)




State, in terms of f , the equation of the graph shown with broken lines.

[1]

.....

(b)




State, in terms of f , the equation of the graph shown with broken lines.

[1]

.....

(c)



State, in terms of f , the equation of the graph shown with broken lines.

[2]

.....

- 4 (a) Expand $(1 + a)^5$ in ascending powers of a up to and including the term in a^3 . [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence expand $[1 + (x + x^2)]^5$ in ascending powers of x up to and including the term in x^3 , simplifying your answer. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is now given instead that the progression is arithmetic.

- (b) (i) Find the common difference of the progression in terms of $\sin \theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Find the sum of the first 16 terms when $\theta = \frac{1}{3}\pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 The functions f and g are defined by

$$f(x) = x^2 - 4x + 3 \quad \text{for } x > c, \text{ where } c \text{ is a constant,}$$

$$g(x) = \frac{1}{x+1} \quad \text{for } x > -1.$$

(a) Express $f(x)$ in the form $(x - a)^2 + b$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is given that f is a one-one function.

(b) State the smallest possible value of c . [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

