

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**FURTHER MATHEMATICS**

**9231/22**

Paper 2

**May/June 2018**

**3 hours**

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF10)

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

Where a numerical value is necessary, take the acceleration due to gravity to be  $10 \text{ m s}^{-2}$ .

The use of a calculator is expected, where appropriate.

Results obtained solely from a graphic calculator, without supporting working or reasoning, will not receive credit.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **22** printed pages and **2** blank pages.



- 1 A bullet of mass  $m$  kg is fired horizontally into a fixed vertical block of material. It enters the block horizontally with speed  $250 \text{ m s}^{-1}$  and emerges horizontally with speed  $70 \text{ m s}^{-1}$  after  $0.04$  s. The block offers a constant horizontal resisting force of magnitude  $450 \text{ N}$ . Find the value of  $m$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 A particle  $P$  moves on a straight line in simple harmonic motion. The centre of the motion is  $O$ . The points  $A$  and  $B$  are on the line, on opposite sides of  $O$ , with  $OA = 1.6 \text{ m}$  and  $OB = 1.2 \text{ m}$ . The ratio of the speed of  $P$  at  $A$  to its speed at  $B$  is  $3 : 4$ .

- (i) Find the amplitude of the motion. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The maximum speed of  $P$  during its motion is  $\frac{1}{3}\pi \text{ m s}^{-1}$ .

(ii) Find the period of the motion. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the time taken for  $P$  to travel directly from  $A$  to  $B$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3** Two identical uniform small spheres  $A$  and  $B$ , each of mass  $m$ , are moving towards each other in a straight line on a smooth horizontal surface. Their speeds are  $u$  and  $ku$  respectively, and they collide directly. The coefficient of restitution between the spheres is  $e$ . Sphere  $B$  is brought to rest by the collision.

**(i)** Show that  $e = \frac{k-1}{k+1}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(ii)** Given that 60% of the total initial kinetic energy is lost in the collision, find the values of  $k$  and  $e$ . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of horizontal dotted lines for writing, consisting of 28 lines.

4 A uniform rod  $AB$  has length  $2a$  and weight  $W$ . The end  $A$  rests on rough horizontal ground and the end  $B$  rests against a smooth vertical wall. The rod is in a vertical plane that is perpendicular to the wall. The angle between the rod and the horizontal is  $\theta$ . A particle of weight  $5W$  hangs from the rod at the point  $C$ , with  $AC = xa$ , where  $0 < x < 1$ .

(i) By taking moments about  $A$ , show that the magnitude of the normal reaction at  $B$  is  $\frac{W(5x + 1)}{2 \tan \theta}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The particle of weight  $5W$  is now moved a distance  $a$  up the rod, so that  $AC = (x + 1)a$ . This results in the magnitude of the normal reaction at  $B$  being double its previous value. The system remains in equilibrium with the rod at angle  $\theta$  with the horizontal.

(ii) Show that  $x = \frac{4}{5}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....





Three thin uniform rings  $A$ ,  $B$  and  $C$  are joined together, so that each ring is in contact with each of the other two rings. Ring  $A$  has radius  $2a$  and mass  $3M$ ; rings  $B$  and  $C$  each have radius  $3a$  and mass  $2M$ . The rings lie in the same plane and the centres of the rings are at the vertices of an isosceles triangle. The object consisting of the three rings is free to rotate about the horizontal axis  $l$  which is tangential to ring  $A$ , in the plane of the object and perpendicular to the line of symmetry of the object (see diagram).

- (i) Show that the moment of inertia of the object about the axis  $l$  is  $180Ma^2$ . [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Show that small oscillations of the object about the axis  $l$  are approximately simple harmonic, and state the period. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 The continuous random variable  $X$  has distribution function given by

$$F(x) = \begin{cases} 1 - e^{-0.4x} & x \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

(i) Find  $P(X > 2)$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the interquartile range of  $X$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- 8 A manufacturer produces three types of car: hatchbacks, saloons and estates. Each type of car is available in one of three colours: silver, blue and red. The manufacturer wants to know whether the popularity of the colour of the car is related to the type of car. A random sample of 300 cars chosen by customers gives the information summarised in the following table.

		Colour of car		
		Silver	Blue	Red
Type of car	Hatchback	53	36	41
	Saloon	29	40	31
	Estate	28	24	18

Test at the 10% significance level whether the colour of car chosen by customers is independent of the type of car. [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of 25 horizontal dotted lines spanning the width of the page, providing a template for writing.

9 At a ski resort, the probability of snow on any particular day is constant and equal to  $p$ . The skiing season begins on 1 November. The random variable  $X$  denotes the day of the skiing season on which the first snowfall occurs. (For example, if the first snowfall is on 5 November, then  $X = 5$ .) The variance of  $X$  is  $\frac{4}{9}$ .

(i) Show that  $4p^2 + 9p - 9 = 0$  and hence find the value of  $p$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the probability that the first snowfall will be on 3 November. [1]

.....

.....

.....

.....

.....

.....

(iii) Find the probability that the first snowfall will not be before 4 November. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iv) Find the least integer  $N$  so that the probability of the first snowfall being on or before the  $N$ th day of November is more than 0.999. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 10 The times taken to run 400 metres by students at two large colleges  $P$  and  $Q$  are being compared. There is no evidence that the population variances are equal. The time taken by a student at college  $P$  and the time taken by a student at college  $Q$  are denoted by  $x$  seconds and  $y$  seconds respectively. A random sample of 50 students from college  $P$  and a random sample of 60 students from college  $Q$  give the following summarised data.

$$\Sigma x = 2620 \quad \Sigma x^2 = 138\,200 \quad \Sigma y = 3060 \quad \Sigma y^2 = 157\,000$$

- (i) Using a 10% significance level, test whether, on average, students from college  $P$  take longer to run 400 metres than students from college  $Q$ . [9]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(ii)** Find a 90% confidence interval for the difference in the mean times taken to run 400 metres by students from colleges *P* and *Q*. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 Answer only **one** of the following two alternatives.

**EITHER**

A particle  $P$  of mass  $m$  is attached to one end of a light inextensible string of length  $a$ . The other end of the string is attached to a fixed point  $O$ . The particle is held so that the string is taut, with  $OP$  horizontal. The particle is projected downwards with speed  $\sqrt{\left(\frac{2}{5}ag\right)}$  and begins to move in a vertical circle. The string breaks when its tension is equal to  $\frac{11}{5}mg$ .

- (i) Show that the string breaks when  $OP$  makes an angle  $\theta$  with the downward vertical through  $O$ , where  $\cos \theta = \frac{3}{5}$ . Find the speed of  $P$  at this instant. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) For the subsequent motion after the string breaks, find the distance  $OP$  when the particle  $P$  is vertically below  $O$ . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



(ii) Find the product moment correlation coefficient for the data.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Test, at the 5% significance level, whether there is evidence of positive correlation between the variables. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Additional Page**

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**BLANK PAGE**

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.